

***T*-SYMMETRICAL TENSOR DIFFERENTIAL FORMS WITH
LOGARITHMIC POLES ALONG A HYPERSURFACE SECTION**

Peter Brückmann^{1 §}, Patrick Winkert²

^{1,2}Department of Mathematics

Martin-Luther-University

Wittenberg, Halle, 06099, GERMANY

¹e-mail: brueckmann@mathematik.uni-halle.de

²e-mail: patrick.winkert@mathematik.uni-halle.de

Abstract: The aim of this paper is to investigate T -symmetrical tensor differential forms with logarithmic poles on the projective space \mathbb{P}^N and on complete intersections $Y \subset \mathbb{P}^N$. Let $H \subset \mathbb{P}^N, N \geq 2$, be a nonsingular irreducible algebraic hypersurface which implies that $D = H$ is a prime divisor in \mathbb{P}^N . The main goal of this paper is the study of the locally free sheaves $\Omega_{\mathbb{P}^N}^T(\log D)$ and the calculation of their cohomology groups. In addition, for complete intersections $Y \subset \mathbb{P}^N$ we give some vanishing theorems and recursion formulas.

AMS Subject Classification: 14F10, 14M10, 14F17, 55N30

Key Words: Young tableaux, complete intersections, algebraic differential forms

1. Introduction

The symmetry properties of tensors are important in physics and certain areas of mathematics. In the following, let k be the ground field which is assumed to be algebraically closed satisfying $\text{char}(k) = 0$. We denote by $H \subset \mathbb{P}_k^N, N \geq 2$, a nonsingular, irreducible, algebraic hypersurface defined by the equation $F = 0$, where $\deg F = m$. Then $D = H$ gives a prime divisor of degree m in \mathbb{P}_k^N . The aim of this paper is the calculation of the dimension of the cohomology groups $H^q(\mathbb{P}^N, \Omega_{\mathbb{P}^N}^T(\log D)(t))$ with general twist $t \in \mathbb{Z}$, where T is a Young

Received: April 25, 2008

© 2008, Academic Publications Ltd.

§Correspondence author

tableau specified later. $\Omega_{\mathbb{P}^N}^T(\log D)$ denotes the so-called sheaf of germs of T -symmetrical tensor differential forms with logarithmic poles along the prime divisor D (cf. [4], [7], [3]). In addition, we consider the associated cohomology groups of nonsingular, irreducible, n -dimensional complete intersections $Y \subset \mathbb{P}^N$, $n \geq 2$. In this case, let the prime divisor $D = Y \cap H$ be the intersection of Y and hypersurface H . As special cases, we investigate the alternating and the symmetric differential forms on \mathbb{P}^N and on Y , respectively.

2. Notations and Preliminaries

Let Ω_X^1 be the sheaf of germs of regular algebraic differential forms on a n -dimensional nonsingular, projective variety $X \subseteq \mathbb{P}^N$ and let $\Omega_X^r = \wedge^r \Omega_X^1$ and $S^r \Omega_X^1$ be the sheaves of alternating and symmetric differential forms on X , alternatively. We denote by $(\Omega_X^1)^{\otimes r}$ the r -th tensor power of Ω_X^1 . The coherent sheaves Ω_X^1 , Ω_X^r , $S^r \Omega_X^1$ and $(\Omega_X^1)^{\otimes r}$ are locally free on X with the rank n , $\binom{n}{r}$, $\binom{n+r-1}{r}$ and n^r , respectively.

The irreducible representations of the symmetric group S_r correspond to the conjugacy classes of S_r . These are given by partitions $(l) : r = l_1 + \dots + l_d$ with $l_i \in \mathbb{Z}$, $l_1 \geq l_2 \geq \dots \geq l_d \geq 1$. Partition (l) can be described by a so-called Young diagram T with r boxes and the row lengths l_1, \dots, l_d . The column lengths of T will be denoted by d_1, \dots, d_l and we set $d = d_1 = \text{depth } T$ and $l = l_1 = \text{length } T$, respectively. Clearly, $d_1 \geq d_2 \geq \dots \geq d_l \geq 1$ and the equations $\sum_{j=1}^l d_j = \sum_{i=1}^d l_i = r$ are fulfilled. Moreover, we put $l_i = 0$ for $i > d$ and $d_j = 0$ for $j > l$. The ‘‘hook-length’’ of the box inside the i -th row and the j -th column of the Young diagram is defined by $h_{i,j} = l_i - i + d_j - j + 1$ and the degree of the associated irreducible representation is equal to

$$\begin{aligned} \nu_{(l)} &= \frac{r!}{\prod h_{i,j}} = \frac{r!}{d!} \cdot \prod_{i=1}^d \frac{i!}{(l_i + d - i)!} \cdot \prod_{1 \leq i < j \leq d} \left(\frac{l_i - l_j}{j - i} + 1 \right) \\ &= r! \cdot \det \left(\left(\frac{1}{\Gamma(l_i + 1 - i + j)} \right)_{i,j=1,\dots,d} \right) \end{aligned}$$

(cf. [5]).

A numbering of the r boxes of a given Young diagram by the integers $1, 2, \dots, r$ in any order is said to be a Young tableau which for simplicity again will be denoted by T . Now, one has an idempotent e_T in the group algebra

$k \cdot S_r$ defined by

$$e_T = \frac{\nu(l)}{r!} \cdot \left(\sum_{q \in Q_T} \text{sgn}(q) \cdot q \right) \circ \left(\sum_{p \in P_T} p \right),$$

where the subgroups P_T and Q_T of S_r are given as follows: $P_T = \{p \in S_r : p \text{ preserves each row of } T\}$, $Q_T = \{q \in S_r : q \text{ preserves each column of } T\}$.

The idempotent e_T is called Young symmetrizer (cf. [5]). If the numbering of the boxes of the Young tableau generates inside every row and every column monotone increasing sequences, we speak of a standard tableau. The number of all standard tableaux to a given Young diagram is equal to the degree $\nu(l)$. We denote by $D(r)$ the set of all standard tableaux to all Young diagrams with r boxes.

For a variety X the notation $\Omega_X^{\otimes r} = (\Omega_X^1)^{\otimes r}$ stands for the sheaf of germs of regular algebraic tensor differential forms. This implies that the symmetric group S_r and the related group algebra $k \cdot S_r$ act on $\Omega_X^{\otimes r}$ defined by $p(a_1 \otimes \dots \otimes a_r) = a_{p^{-1}(1)} \otimes \dots \otimes a_{p^{-1}(r)}$ for all $p \in S_r$. That means, mapping p permutes the spots inside the tensor product. Furthermore, it holds

$$\Omega_X^{\otimes r} = \bigoplus_{T \in D(r)} \Omega_X^T$$

with $\Omega_X^T = e_T(\Omega_X^{\otimes r})$, where Ω_X^T is called the sheaf of germs of T -symmetrical tensor differential forms or simply the T -power of Ω_X^1 . If two Young tableaux T and \tilde{T} possess the same Young diagram, we have $\Omega_X^T \cong \Omega_X^{\tilde{T}}$.

Under the assumption $\text{depth } T \leq \dim X$ with a smooth n -dimensional variety X the belonging sheaf Ω_X^T is locally free of rank

$$\prod_{1 \leq i < j \leq n} \left(\frac{l_i - l_j}{j - i} + 1 \right) = \left(\prod_{i=1}^{n-1} i! \right)^{-1} \cdot \Delta(l_1 - 1, l_2 - 2, \dots, l_n - n),$$

where $\Delta(t_1, t_2, \dots, t_n) = \prod_{1 \leq i < j \leq n} (t_i - t_j)$ denotes the Vandermonde determinant. If $\text{depth } T > \dim X$ then we have $\Omega_X^T = 0$. In the special cases $\Omega_X^r = \wedge^r \Omega_X^1$ and $S^r \Omega_X^1$ the Young tableau has only one column and one row, respectively. In the same way the T -power \mathcal{F}^T of an arbitrary coherent algebraic sheaf \mathcal{F} is defined. One has for instance $\Omega_X^T(\log D) = (\Omega_X^1(\log D))^T$.

Furthermore, we describe the T -power of an algebraic complex (cf. [3]): Let R be a commutative ring which contains the algebraically closed ground field k fulfilling $\text{char}(k) = 0$. We consider an algebraic complex K of R -modules given by $K : K_0 \xrightarrow{d} K_1 \xrightarrow{d} K_2 \xrightarrow{d} \dots$ with $d^2 = 0$.

Then the r -th tensor power $P = K^{\otimes r}$ of K is defined by $P = K^{\otimes r} : P_0 \xrightarrow{\delta} P_1 \xrightarrow{\delta} P_2 \xrightarrow{\delta} \dots$ with $P_s = \bigoplus_{s_1+\dots+s_r=s} K_{s_1} \otimes \dots \otimes K_{s_r}$ and $\delta(b_1 \otimes \dots \otimes b_r) = \sum_{i=1}^r (-1)^{s_1+\dots+s_{i-1}} \cdot b_1 \otimes \dots \otimes b_{i-1} \otimes d(b_i) \otimes b_{i+1} \otimes \dots \otimes b_r$, where $b_j \in K_{s_j}$ for all j . Again the symmetric group S_r acts on this tensor power by permutation of the spots inside the tensor product. In order to obtain such an action of S_r on $P = K^{\otimes r}$, which commutes with δ , we introduce additionally a sign as follows:

1. $\sigma(p; s_1, \dots, s_r) := \sum_{\substack{i < j \\ p(i) > p(j)}} s_i \cdot s_j$ for all $p \in S_r$.
2. $p(b_1 \otimes \dots \otimes b_r) := (-1)^{\sigma(p; s_1, \dots, s_r)} \cdot b_{p^{-1}(1)} \otimes \dots \otimes b_{p^{-1}(r)}$, where $b_j \in K_{s_j}$ for all $j \in \{1, \dots, r\}$.

Then one has

$$P_s = \bigoplus_{T \in D(r)} K_s^{(T)}, \quad K^{\otimes r} = \bigoplus_{T \in D(r)} K^{(T)}, \quad H^*(K^{\otimes r}) = \bigoplus_{T \in D(r)} H^*(K^{(T)})$$

with $K_s^{(T)} = e_T(P_s)$ and $K^{(T)} = e_T(K^{\otimes r}) : K_0^{(T)} \xrightarrow{\delta} K_1^{(T)} \xrightarrow{\delta} K_2^{(T)} \xrightarrow{\delta} \dots$

This complex $K^{(T)}$ is said to be the T -power of K . If two Young tableaux T and \tilde{T} possess the same Young diagram, one has $K^{(T)} \cong K^{(\tilde{T})}$. For an exact sequence K the T -power $K^{(T)}$ of K is also an exact sequence.

Now, let $X \subseteq \mathbb{P}^N$ be a projective variety satisfying $\omega_X \cong \mathcal{O}_X(n_X)$ for some $n_X \in \mathbb{Z}$, where ω_X stands for the canonical line bundle. This implies under the assumptions $d = \text{depth } T = \dim X = n$ and $l = \text{length } T > 1$ the isomorphism

$$\Omega_X^T \cong \Omega_X^{T'} \otimes \omega_X \cong \Omega_X^{T'}(n_X),$$

where T' arises from T by deleting the first column of T . In the case $d = \text{depth } T = \dim X = n$ and $l = \text{length } T = 1$ (i.e. T has only one column) we have the isomorphism $\Omega_X^T \cong \Omega_X^n \cong \omega_X \cong \mathcal{O}_X(n_X)$.

An important tool in our considerations will be the Serre duality: Suppose the Young tableau T has the column lengths d_1, \dots, d_l satisfying $d_1 = d = \text{depth } T \leq \dim X = n$. We get an associated Young tableau T^* by the column lengths $d_j^* = n - d_{l+1-j}$ for all $j = 1, \dots, l$. One verifies readily that in case $\text{depth } T < n$ holds $(T^*)^* = T$.

The next lemma delivers some duality relations about the dimensions of cohomology groups.

Lemma 2.1. *Let $Y = H_1 \cap \dots \cap H_{N-n} \subseteq \mathbb{P}^N$ be a n -dimensional, non-singular, irreducible, complete intersection defined by algebraic hypersurfaces $H_i \subset \mathbb{P}^N$ satisfying $F_i = 0$ with $\deg F_i = m_i$. The dimension of Y is n . In this case, let the prime divisor $D = Y \cap H$ be the intersection of Y and hypersurface*

$H : F = 0$ of degree m . Assume that D also becomes a nonsingular irreducible complete intersection of dimension $n - 1$. Then one has:

- (i) $\dim H^q(\mathbb{P}^N, \Omega_{\mathbb{P}^N}^r(\log D)(t)) = \dim H^{N-q}(\mathbb{P}^N, \Omega_{\mathbb{P}^N}^{N-r}(\log D)(-t - m))$.
- (ii) $\dim H^q(Y, \Omega_Y^r(\log D)(t)) = \dim H^{n-q}(Y, \Omega_Y^{n-r}(\log D)(-t - m))$.
- (iii) $\dim H^q(\mathbb{P}^N, \Omega_{\mathbb{P}^N}^T(\log D)(t)) = \dim H^{N-q}(\mathbb{P}^N, \Omega_{\mathbb{P}^N}^{T^*}(\log D)(-t - l \cdot m + (l - 1)(N + 1)))$.
- (iv) $\dim H^q(Y, \Omega_Y^T(\log D)(t)) = \dim H^{n-q}(Y, \Omega_Y^{T^*}(\log D)(-t - l \cdot m - (l - N - n) \cdot 1))$.
- (v) $\dim H^q(\mathbb{P}^N, S^r \Omega_{\mathbb{P}^N}^1(\log D)(t)) = \dim H^{N-q}(\mathbb{P}^N, \Omega_{\mathbb{P}^N}^{T^*}(\log D)(-t - r \cdot m + (r - 1)(N + 1)))$, where T^* denotes a rectangle with $N - 1$ rows and r columns.
- (vi) $\dim H^q(Y, S^r \Omega_Y^1(\log D)(t)) = \dim H^{n-q}(Y, \Omega_Y^{T^*}(\log D)(-t - r \cdot m - (r - N - n) \cdot 1))$, where T^* denotes a rectangle with $n - 1$ rows and r columns.

Proof. We consider the following exact sequence (cf. [4])

$$0 \longrightarrow \Omega_{\mathbb{P}^N}^r(\log D)(-m) \longrightarrow \Omega_{\mathbb{P}^N}^r \longrightarrow \Omega_D^r \longrightarrow 0.$$

For $r = N$ we have $\Omega_D^N = 0$, i.e. $\Omega_{\mathbb{P}^N}^N(\log D) \cong \Omega_{\mathbb{P}^N}^N(m) \cong \mathcal{O}_{\mathbb{P}^N}(m - N - 1)$. This implies a pairing $\Omega_{\mathbb{P}^N}^r(\log D)(t) \times \Omega_{\mathbb{P}^N}^{N-r}(\log D)(-t - m + N + 1) \longrightarrow \mathcal{O}_{\mathbb{P}^N}$, which means that the vector space $H^q(\mathbb{P}^N, \Omega_{\mathbb{P}^N}^r(\log D)(t))$ is dual to $H^{N-q}(\mathbb{P}^N, (\Omega_{\mathbb{P}^N}^{N-r}(\log D)(-t - m + N + 1)) \otimes \Omega_{\mathbb{P}^N}^N)$.

Setting $\Omega_{\mathbb{P}^N}^N \cong \mathcal{O}_{\mathbb{P}^N}(-N - 1)$ yields (i). The statement (ii) can be shown in a similar way. Note that $\Omega_Y^n(\log D) \cong \Omega_Y^n(m) \cong \mathcal{O}_Y(m + \sum_{i=1}^{N-n} m_i - N - 1)$. Now, let T be a Young tableau with r boxes, given by the row lengths l_1, \dots, l_d and the column lengths d_1, \dots, d_l where $d = d_1 = \text{depth } T$ and $l = l_1 = \text{length } T$. The Young tableau T^* has the column lengths $d_j^* = n - d_{l+1-j}$ for all $j \in \{1, \dots, l\}$ and we have again $\Omega_{\mathbb{P}^N}^N(\log D) \cong \mathcal{O}_{\mathbb{P}^N}(m - N - 1)$. From the pairing $\Omega_{\mathbb{P}^N}^T(\log D)(t) \times \Omega_{\mathbb{P}^N}^{T^*}(\log D)(-t - l \cdot (m - N - 1)) \longrightarrow \mathcal{O}_{\mathbb{P}^N}$ follows $\text{Hom}(\Omega_{\mathbb{P}^N}^T(\log D)(t), \mathcal{O}_{\mathbb{P}^N}) \cong \Omega_{\mathbb{P}^N}^{T^*}(\log D)(-t - l \cdot (m - N - 1))$, which shows assertion (iii). In order to show the formula for complete intersections Y instead of \mathbb{P}^N , we replace $-N - 1$ by $\sum_{i=1}^{N-n} m_i - N - 1$. Choosing $l = r$ (depth $T = 1$) in (iii) and (iv) proves (v) and (vi), respectively. \square

For a projective variety $X \subseteq \mathbb{P}^N$ and a coherent sheaf \mathcal{F} on X the dimensions $\dim_k H^q(X, \mathcal{F})$ are finite and we have the so-called Euler-Poincaré

characteristic given by $\chi(X, \mathcal{F}) = \sum_{q=0}^{\dim X} (-1)^q \cdot \dim H^q(X, \mathcal{F})$. From a short exact sequence $0 \rightarrow \mathcal{F} \rightarrow \mathcal{G} \rightarrow \mathcal{H} \rightarrow 0$ with coherent sheaves $\mathcal{F}, \mathcal{G}, \mathcal{H}$ on X we obtain the equation $\chi(X, \mathcal{G}) = \chi(X, \mathcal{F}) + \chi(X, \mathcal{H})$. Under the above assumptions we also know, that for a short exact sequence of coherent sheaves on X there exists a long exact sequence for the associated cohomology groups. For every coherent sheaf \mathcal{F} on the projective variety $X \subset \mathbb{P}^N$ there exists a polynomial $P(X, \mathcal{F})(t) \in \mathbb{Q}[t]$ of degree $\dim X$ which fulfills $\chi(X, \mathcal{F}(t)) = P(X, \mathcal{F})(t)$ for all $t \in \mathbb{Z}$. $P(X, \mathcal{F})(t)$ is said to be the Hilbert polynomial of \mathcal{F} (cf. [8], [6], [7]). For example, the structure sheaf on \mathbb{P}^N has the following Hilbert polynomial

$$P(\mathbb{P}^N, \mathcal{O}_{\mathbb{P}^N})(t) = \chi(\mathbb{P}^N, \mathcal{O}_{\mathbb{P}^N}(t)) = \frac{(t+N) \cdot \dots \cdot (t+1)}{N!}. \quad (2.1)$$

3. The Projective Space \mathbb{P}^N

In the following, we change the meaning of the binomial coefficient setting $\binom{\alpha}{\beta} = 0$ for all $\alpha \in \mathbb{Z}, \beta \in \mathbb{N}$ satisfying $\alpha < \beta$, in particular: $\binom{\alpha}{\beta} = 0$ if $\alpha < 0$. For instance: $\dim H^0(\mathbb{P}^N, \mathcal{O}_{\mathbb{P}^N}(t)) = \binom{t+N}{N}$, $\dim H^N(\mathbb{P}^N, \mathcal{O}_{\mathbb{P}^N}(t)) = \binom{-t-1}{N}$, $H^q(\mathbb{P}^N, \mathcal{O}_{\mathbb{P}^N}(t)) = 0$ for $0 < q < N$.

Let $H \subset \mathbb{P}^N$ ($N \geq 2$) be a nonsingular, irreducible, algebraic hypersurface defined by the equation $F = 0$, that means, $D = H$ is a prime divisor in \mathbb{P}^N . Both F and D are of degree m and $D = H$ has dimension $N - 1$.

3.1. Alternating Differential Forms

We denote by $\Omega_{\mathbb{P}^N}^r$ the local free sheaf of germs of alternating differential forms on the projective space \mathbb{P}^N and consider the following sequence ($t \in \mathbb{Z}$)

$$0 \longrightarrow \Omega_{\mathbb{P}^N}^r(t) \longrightarrow \Omega_{\mathbb{P}^N}^r(\log D)(t) \longrightarrow \Omega_D^{r-1}(t) \longrightarrow 0, \quad (3.1)$$

which is known to be exact (cf. [4]). The dimensions of the cohomology groups $H^q(\mathbb{P}^N, \Omega_{\mathbb{P}^N}^r(t))$ and $H^q(D, \Omega_D^{r-1}(t))$ are calculated in [1], where we also find the following exact sequences

$$0 \longrightarrow \Omega_{\mathbb{P}^N}^r(t-m) \longrightarrow \Omega_{\mathbb{P}^N}^r(t) \xrightarrow{\alpha} \mathcal{O}_D(t) \otimes_{\mathcal{O}_{\mathbb{P}^N}} \Omega_{\mathbb{P}^N}^r \longrightarrow 0, \quad (3.2)$$

$$0 \longrightarrow \Omega_D^{r-1}(t-m) \longrightarrow \mathcal{O}_D(t) \otimes_{\mathcal{O}_{\mathbb{P}^N}} \Omega_{\mathbb{P}^N}^r \xrightarrow{\beta} \Omega_D^r(t) \longrightarrow 0. \quad (3.3)$$

The mapping $\varphi^* := \beta \circ \alpha$ means the restriction of the differential forms on \mathbb{P}^N to the hypersurface $D = H$. In the case $r = 1$, one has to replace the sheaf Ω_D^{r-1} by

the structure sheaf \mathcal{O}_D . For $0 < q < N$ we have $\dim H^q(\mathbb{P}^N, \Omega_{\mathbb{P}^N}^r(t)) = \delta_{q,r} \cdot \delta_{t,0}$ (Kronecker- δ) and we know by [1, Lemma 4] a base element of $H^r(\mathbb{P}^N, \Omega_{\mathbb{P}^N}^r)$ which is given by the cohomology class of the cocycle $\omega^{(r)} \in C^r(\mathfrak{U}, \Omega_{\mathbb{P}^N}^r)$ defined by

$$\omega_{i_0, \dots, i_r}^{(r)} = \frac{x_{i_0}}{x_{i_r}} \cdot d \frac{x_{i_1}}{x_{i_0}} \wedge d \frac{x_{i_2}}{x_{i_1}} \wedge \dots \wedge d \frac{x_{i_r}}{x_{i_{r-1}}}. \tag{3.4}$$

\mathfrak{U} stands for the affine open covering of \mathbb{P}^N by the affine spaces $U_i = \{x_i \neq 0\}$. For $r = 1$, in particular, $\omega_{i_0, i_1}^{(1)} = \frac{x_{i_0}}{x_{i_1}} \cdot d \frac{x_{i_1}}{x_{i_0}}$ is a logarithmic differential. We may represent (3.4) by

$$\omega_{i_0, \dots, i_r}^{(r)} = \omega_{i_0, i_1}^{(1)} \wedge \omega_{i_1, i_2}^{(1)} \wedge \dots \wedge \omega_{i_{r-1}, i_r}^{(1)},$$

which is an outer product of logarithmic differential forms. In the case $q = r = N$, $t = 0$ the cochain $\omega^{(N)}$ creates a base of $H^N(\mathbb{P}^N, \Omega_{\mathbb{P}^N}^N)$ (cf. [1, Lemma 2]). Finally, we set $\omega^{(0)} = 1$.

Lemma 3.1. *Let $0 < r \leq N$. Then the homomorphism $d : H^{r-1}(D, \Omega_D^{r-1}) \rightarrow H^r(\mathbb{P}^N, \Omega_{\mathbb{P}^N}^r)$ in the long homology sequence with respect to the exact sequence*

$$0 \rightarrow \Omega_{\mathbb{P}^N}^r \rightarrow \Omega_{\mathbb{P}^N}^r(\log D) \rightarrow \Omega_D^{r-1} \rightarrow 0$$

is epimorphic. If in addition $2(r-1) \neq N-1$ is valid, then d is an isomorphism.

Proof. We calculate the image of the cohomology class of $\omega^{(r-1)}$ at the composition

$$H^{r-1}(\mathbb{P}^N, \Omega_{\mathbb{P}^N}^{r-1}) \xrightarrow{\varphi^*} H^{r-1}(D, \Omega_D^{r-1}) \xrightarrow{d} H^r(\mathbb{P}^N, \Omega_{\mathbb{P}^N}^r)$$

and denote $\varphi^*(\omega^{(r-1)})$ again by $\omega^{(r-1)}$. Let \mathfrak{U} be the affine, open covering of \mathbb{P}^N given by the affine spaces $U_i = \{x_i \neq 0\}$. We consider the following commutative diagram

$$\begin{array}{ccccccc} 0 & \rightarrow & C^{r-1}(\mathfrak{U}, \Omega_{\mathbb{P}^N}^r) & \rightarrow & C^{r-1}(\mathfrak{U}, \Omega_{\mathbb{P}^N}^r(\log D)) & \rightarrow & C^{r-1}(\mathfrak{U}, \Omega_D^{r-1}) \rightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \rightarrow & C^r(\mathfrak{U}, \Omega_{\mathbb{P}^N}^r) & \rightarrow & C^r(\mathfrak{U}, \Omega_{\mathbb{P}^N}^r(\log D)) & \rightarrow & C^r(\mathfrak{U}, \Omega_D^{r-1}) \rightarrow 0 \end{array}$$

where the cocycle $\omega^{(r-1)} \in C^{r-1}(\mathfrak{U}, \Omega_D^{r-1})$ possesses in $C^{r-1}(\mathfrak{U}, \Omega_{\mathbb{P}^N}^r(\log D))$ the preimage ϱ defined by $\varrho_{i_0, \dots, i_{r-1}} = \omega_{i_0, \dots, i_{r-1}}^{(r-1)} \wedge \frac{x_{i_0}^m}{F^r} \cdot d \frac{F}{x_{i_0}^m}$ (cf. [4]).

Elementary calculations show that $d\omega^{(r-1)} = (-1)^r \cdot m \cdot \omega^{(r)} \in C^r(\mathfrak{U}, \Omega_{\mathbb{P}^N}^r)$. Therefore, the cocycle $d\omega^{(r-1)} \in C^r(\mathfrak{U}, \Omega_{\mathbb{P}^N}^r)$ is nonzero and the associated cohomology class is a base of $H^r(\mathbb{P}^N, \Omega_{\mathbb{P}^N}^r)$. Thus, the homomorphism $d : H^{r-1}(D, \Omega_D^{r-1}) \rightarrow H^r(\mathbb{P}^N, \Omega_{\mathbb{P}^N}^r)$ is epimorphic. In the case $2(r-1) \neq N-1$,

we obtain $\dim H^{r-1}(D, \Omega_D^{r-1}) = 1$ by [1, Satz 2 and Lemma 5], which implies that d is an isomorphism. \square

Theorem 3.2. *Let $D \subset \mathbb{P}^N$ be a smooth algebraic hypersurface of degree m ($N \geq 2$).*

(a) *For each $r \in \{1, \dots, N-1\}$ one has:*

$$\dim H^0(\mathbb{P}^N, \Omega_{\mathbb{P}^N}^r(\log D)(t)) = \sum_{i=0}^r (-1)^i \cdot \binom{N+1}{r-i} \cdot \binom{t+N-i \cdot (m-1) - r}{N}.$$

(b) *For all $r \in \{1, \dots, N-1\}$ it holds: $H^0(\mathbb{P}^N, \Omega_{\mathbb{P}^N}^r(\log D)(t)) \neq 0 \Leftrightarrow t \geq r$.*

(c) *In the case $r = N$ one has: $\dim H^0(\mathbb{P}^N, \Omega_{\mathbb{P}^N}^N(\log D)(t)) = \binom{t+m-1}{N}$.*

(d) *If $D \subset \mathbb{P}^N$ is a hyperplane ($m = 1$), then it holds: $\dim H^0(\mathbb{P}^N, \Omega_{\mathbb{P}^N}^r(\log D)(t)) = \binom{N}{r} \cdot \binom{t+N-r}{N}$.*

Proof. The formula (a) follows directly from the long exact cohomology sequence related to the exact sequence in (3.1) by applying Lemma 3.1. For $r = N$ we obtain $\Omega_{\mathbb{P}^N}^N(\log D) \cong \Omega_{\mathbb{P}^N}^N(m) \cong \mathcal{O}_{\mathbb{P}^N}(m - N - 1)$ which yields (c). (a) obviously implies (b) and (d). \square

Theorem 3.3. (a) *Let $0 < q < N$, $q + r \neq N$ and $r \geq 1$. Then we obtain $H^q(\mathbb{P}^N, \Omega_{\mathbb{P}^N}^r(\log D)(t)) = 0$ for all $t \in \mathbb{Z}$.*

(b) *For $1 \leq r \leq N-1$ it follows:*

$$\begin{aligned} & \dim H^{N-r}(\mathbb{P}^N, \Omega_{\mathbb{P}^N}^r(\log D)(t)) \\ &= \sum_{i=0}^{N+1} (-1)^i \cdot \binom{N+1}{i} \cdot \binom{t + (N-r) \cdot m - (i-1) \cdot (m-1)}{N} \\ &= \sum_{i=0}^{N+1} (-1)^i \cdot \binom{N+1}{i} \cdot \binom{-t + (r-1) \cdot m - (i-1) \cdot (m-1)}{N} \end{aligned}$$

That means: If D is a hyperplane ($m = 1$), then we have $H^{N-r}(\mathbb{P}^N, \Omega_{\mathbb{P}^N}^r(\log D)(t)) = 0$ for all $t \in \mathbb{Z}$.

(c) *For $1 \leq r \leq N-1$ one has:*

$$\begin{aligned} & \dim H^N(\mathbb{P}^N, \Omega_{\mathbb{P}^N}^r(\log D)(t)) \\ &= \sum_{i=0}^{N-r} (-1)^i \cdot \binom{N+1}{N-r-i} \cdot \binom{-t - m - i \cdot (m-1) + r}{N} \end{aligned}$$

If D is a hyperplane ($m = 1$), then we get: $\dim H^N(\mathbb{P}^N, \Omega_{\mathbb{P}^N}^r(\log D)(t)) = \binom{N}{r} \cdot \binom{-t-1+r}{N}$.

$$(d) \dim H^N(\mathbb{P}^N, \Omega_{\mathbb{P}^N}^N(\log D)(t)) = \binom{-t-m+N}{N}.$$

Proof. We consider the following exact sequence

$$\begin{aligned} \dots &\longrightarrow H^{q-1}(D, \Omega_D^{r-1}(t)) \xrightarrow{d_1} H^q(\mathbb{P}^N, \Omega_{\mathbb{P}^N}^r(t)) \longrightarrow \\ &\longrightarrow H^q(\mathbb{P}^N, \Omega_{\mathbb{P}^N}^r(\log D)(t)) \longrightarrow H^q(D, \Omega_D^{r-1}(t)) \xrightarrow{d_2} \\ &\xrightarrow{d_2} H^{q+1}(\mathbb{P}^N, \Omega_{\mathbb{P}^N}^r(t)) \longrightarrow \dots, \end{aligned} \tag{3.5}$$

and assume $0 < q < r$ and $q + r < N$. By Lemma 3.1 the mappings d_1 and d_2 are epimorphic for all $t \in \mathbb{Z}$ and from (3.5) we get the exact sequence

$$0 \rightarrow H^q(\mathbb{P}^N, \Omega_{\mathbb{P}^N}^r(\log D)(t)) \rightarrow H^q(D, \Omega_D^{r-1}(t)) \xrightarrow{d_2} H^{q+1}(\mathbb{P}^N, \Omega_{\mathbb{P}^N}^r(t)) \rightarrow 0.$$

Under these assumptions holds $H^q(D, \Omega_D^{r-1}(t)) = 0$ if $q \neq r - 1$ or $t \neq 0$ (cf. [1]). In case $q = r - 1, t = 0$ we know that d_2 is an isomorphism by Lemma 3.1 since $2(r - 1) < N - 1$. Therefore, one has

$$H^q(\mathbb{P}^N, \Omega_{\mathbb{P}^N}^r(\log D)(t)) = 0 \quad \text{for } 0 < q < r \text{ and } q + r < N.$$

For $q < N, r < N, q + r > N$ we use the Serre duality to show statement (a). The case $r = N$ is trivial since $\Omega_{\mathbb{P}^N}^N(\log D) \cong \mathcal{O}_{\mathbb{P}^N}(m - N - 1)$.

If $r \geq 2$ and $q + r = N$ then the mappings d_1 and d_2 are epimorphic, i.e.

$$\begin{aligned} \dim H^{N-r}(\mathbb{P}^N, \Omega_{\mathbb{P}^N}^r(\log D)(t)) \\ = \dim H^{N-r}(D, \Omega_D^{r-1}(t)) - \dim H^{N-r+1}(\mathbb{P}^N, \Omega_{\mathbb{P}^N}^r(t)). \end{aligned}$$

In the case $r = 1$ and $q = N - 1$ one has $\dim H^{N-1}(\mathbb{P}^N, \Omega_{\mathbb{P}^N}^1(\log D)(t)) = \dim H^{N-1}(D, \mathcal{O}_D(t)) - \dim H^N(\mathbb{P}^N, \Omega_{\mathbb{P}^N}^1(t)) + H^N(\mathbb{P}^N, \Omega_{\mathbb{P}^N}^1(\log D)(t))$.

Applying Theorem 3.2, Lemma 2.1 and the results in [1] delivers (b) and (c). □

3.2. T-Symmetric Tensor Differential Forms

Let T be a Young tableau with r boxes. We study the sheaf $\Omega^T(\log D) = (\Omega^1(\log D))^T$ on \mathbb{P}^N and begin with a free resolution of the sheaf $\Omega^1(\log D)$.

Lemma 3.4. *Let $D \subset \mathbb{P}^N$ be a nonsingular, irreducible, algebraic hypersurface of degree $m \geq 2$ defined by the equation $F = 0$.*

Then there exists a short exact sequence

$$0 \longrightarrow \mathcal{O}_{\mathbb{P}^N}(-m) \longrightarrow \bigoplus^{N+1} \mathcal{O}_{\mathbb{P}^N}(-1) \longrightarrow \Omega_{\mathbb{P}^N}^1(\log D) \longrightarrow 0. \quad (3.6)$$

If D is a hyperplane, i.e. $m = 1$, we have $\Omega_{\mathbb{P}^N}^1(\log D) \cong \bigoplus^N \mathcal{O}_{\mathbb{P}^N}(-1)$.

Proof. Let $U_i = \{x_i \neq 0\} \subset \mathbb{P}^N$ and let $U \subseteq \mathbb{P}^N$ be an arbitrary open affine subset. We are going to show that there is an exact sequence

$$0 \rightarrow \Gamma(U, \mathcal{O}_{\mathbb{P}^N}(-m)) \xrightarrow{\alpha} \bigoplus^{N+1} \Gamma(U, \mathcal{O}_{\mathbb{P}^N}(-1)) \xrightarrow{\beta} \Gamma(U, \Omega_{\mathbb{P}^N}^1(\log D)) \rightarrow 0.$$

For sections $f_0, \dots, f_N \in \Gamma(U, \mathcal{O}(-1))$ we put $g := -\frac{1}{m} \cdot \sum_{\mu=0}^N x_\mu f_\mu \in \Gamma(U, \mathcal{O})$. Let $F_j = \frac{\partial F}{\partial x_j}$ denotes the partial derivatives of F . The mapping β is defined by $(f_0, \dots, f_N) \mapsto \omega$, where the differential form ω on $U \cap U_i$ is given by

$$\omega = \omega_i := \sum_{\substack{\nu=0 \\ \nu \neq i}}^N \left(f_\nu + g \cdot \frac{F_\nu}{F} \right) \cdot x_i \cdot d \frac{x_\nu}{x_i}.$$

One easily verifies that ω is a section of $\Omega_{\mathbb{P}^N}^1(\log D)$ on U and it holds, in particular, $\omega_i = \omega_j$ for any $i, j \in \{0, 1, \dots, N\}$. For a section $\delta \in \Gamma(U, \mathcal{O}(-m))$ let $f_\nu = \delta \cdot F_\nu$ for all $\nu = 0, 1, \dots, N$ which implies that $f_\nu \in \Gamma(U, \mathcal{O}(-1))$ and $g = -\delta \cdot F$. Finally, we have $\ker \beta = \{(\delta \cdot F_0, \dots, \delta \cdot F_N)\} \cong \Gamma(U, \mathcal{O}_{\mathbb{P}^N}(-m))$, which yields the claim for $m \geq 2$.

In the last part we have to show the statement of Lemma 3.4 in case $m = 1$. Let $D \subset \mathbb{P}^N$ be the hyperplane satisfying the equation $x_N = 0$, and let $U \subseteq \mathbb{P}^N$ be an open subset. For given sections $f_0, \dots, f_{N-1} \in \Gamma(U, \mathcal{O}(-1))$ let ω be the differential form, which has on $U \cap U_i$ $i = 0, \dots, N-1$, the representation

$$\omega = \omega_i = \sum_{\substack{\nu=0 \\ \nu \neq i}}^{N-1} f_\nu \cdot x_i \cdot d \frac{x_\nu}{x_i} - \left(\sum_{\mu=0}^{N-1} f_\mu \cdot x_\mu \right) \cdot \frac{x_i}{x_N} d \frac{x_N}{x_i},$$

respectively on $U \cap U_N$,

$$\omega = \omega_N = \sum_{\nu=0}^{N-1} f_\nu \cdot x_N \cdot d \frac{x_\nu}{x_N}.$$

Then ω is a section of $\Omega_{\mathbb{P}^N}^1(\log D)$ on U , and the mapping $(f_0, \dots, f_{N-1}) \mapsto \omega$ becomes an isomorphism of $\Gamma(U, \bigoplus^N \mathcal{O}_{\mathbb{P}^N}(-1))$ onto $\Gamma(U, \Omega_{\mathbb{P}^N}^1(\log D))$. \square

Lemma 3.5. *Let T be a Young tableau with r boxes and the row lengths l_1, l_2, \dots, l_d , set $t_i := r + l_i - i$ for all $i \geq 1$ ($l_i = 0$ if $i > d$) and assume*

$d = \text{depth } T \leq N$. Then the following sequence is exact for $m \geq 2$:

$$\begin{aligned}
 0 \longrightarrow \bigoplus_{b_d} \mathcal{O}_{\mathbb{P}^N}(d \cdot (1 - m) - r) &\xrightarrow{\alpha_d} \bigoplus_{b_{d-1}} \mathcal{O}_{\mathbb{P}^N}((d - 1) \cdot (1 - m) - r) \xrightarrow{\alpha_{d-1}} \\
 &\xrightarrow{\alpha_{d-1}} \bigoplus_{b_{d-2}} \mathcal{O}_{\mathbb{P}^N}((d - 2) \cdot (1 - m) - r) \xrightarrow{\alpha_{d-2}} \dots \xrightarrow{\alpha_2} \bigoplus_{b_1} \mathcal{O}_{\mathbb{P}^N}(1 - m - r) \xrightarrow{\alpha_1} \\
 &\xrightarrow{\alpha_1} \bigoplus_{b_0} \mathcal{O}_{\mathbb{P}^N}(-r) \xrightarrow{\alpha_0} \Omega_{\mathbb{P}^N}^T(\log D) \longrightarrow 0
 \end{aligned}
 \tag{3.7}$$

with the integers

$$b_s = \left(\prod_{i=1}^N i! \right)^{-1} \cdot \sum_{1 \leq i_1 < \dots < i_s \leq d} \Delta(t_1, t_2, \dots, t_{i_1} - 1, \dots, t_{i_s} - 1, \dots, t_N, t_{N+1}), \tag{3.8}$$

where Δ denotes the Vandermonde determinant.

In the case $s = 0$ we have $b_0 = \left(\prod_{i=1}^N i! \right)^{-1} \cdot \prod_{1 \leq i < j \leq N+1} (l_i - l_j + j - i) = \left(\prod_{i=1}^N i! \right)^{-1} \cdot \Delta(t_1, t_2, \dots, t_N, t_{N+1})$. For $m = 1$ (D is a hyperplane) holds

$$\Omega_{\mathbb{P}^N}^T(\log D) \cong \bigoplus^{\text{rk}(\Omega_{\mathbb{P}^N}^T)} \mathcal{O}_{\mathbb{P}^N}(-r)$$

with

$$\begin{aligned}
 \text{rk}(\Omega_{\mathbb{P}^N}^T) &= \prod_{1 \leq i < j \leq N} \left(\frac{l_i - l_j}{j - i} + 1 \right) = \sum_{i=0}^d (-1)^i \cdot b_i \\
 &= \left(\prod_{i=1}^{N-1} i! \right)^{-1} \cdot \Delta(t_1, t_2, \dots, t_N).
 \end{aligned}$$

Proof. The T-Power of (3.6) yields the claim for $m \geq 2$ (cf. [3]) and Lemma 3.4 shows the case $m = 1$. □

Theorem 3.6. Let T be a Young tableau with r boxes and with $d = \text{depth } T$ rows.

$$(a) \chi(\mathbb{P}^N, \Omega_{\mathbb{P}^N}^T(\log D)(t)) = \frac{1}{N!} \cdot \sum_{i=0}^d (-1)^i \cdot b_i \cdot \prod_{j=1}^N (t - i \cdot (m - 1) + j - r).$$

(b) For depth $T < N$ one has:

$$\dim H^0(\mathbb{P}^N, \Omega_{\mathbb{P}^N}^T(\log D)(t)) = \sum_{i=0}^d (-1)^i \cdot b_i \cdot \binom{t - i \cdot (m - 1) + N - r}{N}$$

and therefore: $H^0(\mathbb{P}^N, \Omega_{\mathbb{P}^N}^T(\log D)(t)) \neq 0 \Leftrightarrow t \geq r$.

(c) Let $d = \text{depth } T = N$ and let l_N be the number of columns of T with the length N . We denote by T' the Young tableau which is given by T without these columns of length N . Then $\text{depth } T' < N$ and it holds $\Omega_{\mathbb{P}^N}^T(\log D)(t) \cong \Omega_{\mathbb{P}^N}^{T'}(\log D)(t + l_N \cdot (m - N - 1))$.

If T is a rectangle with N rows and l columns, then we have $\Omega_{\mathbb{P}^N}^T(\log D)(t) \cong \mathcal{O}_{\mathbb{P}^N}(t + l \cdot (m - N - 1))$.

(d) For $1 \leq q < N - d$ we get $H^q(\mathbb{P}^N, \Omega_{\mathbb{P}^N}^T(\log D)(t)) = 0$ for all $t \in \mathbb{Z}$.

(e) Let d_l be the length of the last column of T . Then it holds: $H^q(\mathbb{P}^N, \Omega_{\mathbb{P}^N}^T(\log D)(t)) = 0$ for $N - d_l < q < N$ and $\forall t \in \mathbb{Z}$.

Proof. The short exact sequences of (3.7) yields

$$\begin{array}{ccccccc} 0 & \longrightarrow & \bigoplus_{b_d} \mathcal{O}_{\mathbb{P}^N}(d \cdot (1 - m) - r) & \longrightarrow & \bigoplus_{b_{d-1}} \mathcal{O}_{\mathbb{P}^N}((d - 1) \cdot (1 - m) - r) & \longrightarrow & \\ & & \longrightarrow \text{Im } \alpha_{d-1} & \longrightarrow & 0, & & \\ 0 & \longrightarrow & \text{Im } \alpha_{d-1} & \longrightarrow & \bigoplus_{b_{d-2}} \mathcal{O}_{\mathbb{P}^N}((d - 2) \cdot (1 - m) - r) & \longrightarrow & \text{Im } \alpha_{d-2} \longrightarrow 0, \\ & & & & \vdots & & \vdots \\ 0 & \longrightarrow & \text{Im } \alpha_2 & \longrightarrow & \bigoplus_{b_1} \mathcal{O}_{\mathbb{P}^N}(1 - m - r) & \longrightarrow & \text{Im } \alpha_1 \longrightarrow 0, \\ 0 & \longrightarrow & \text{Im } \alpha_1 & \longrightarrow & \bigoplus_{b_0} \mathcal{O}_{\mathbb{P}^N}(-r) & \longrightarrow & \Omega_{\mathbb{P}^N}^T(\log D) \longrightarrow 0, \end{array}$$

where $H^q(\mathbb{P}^N, \mathcal{O}_{\mathbb{P}^N}(t)) = 0$ for $1 \leq q \leq N - 1$ and for all $t \in \mathbb{Z}$. This implies $H^q(\mathbb{P}^N, \text{Im } \alpha_i(t)) = 0$ for $1 \leq q \leq N - 1 + i - d$ and hence, we have in case $d < N$

$$\begin{aligned} & \dim H^0(\mathbb{P}^N, \Omega_{\mathbb{P}^N}^T(\log D)(t)) \\ &= b_0 \cdot \dim H^0(\mathbb{P}^N, \mathcal{O}_{\mathbb{P}^N}(t - r)) - \dim H^0(\mathbb{P}^N, \text{Im } \alpha_1(t)), \\ & \dim H^0(\mathbb{P}^N, \text{Im } \alpha_1(t)) \\ &= b_1 \cdot \dim H^0(\mathbb{P}^N, \mathcal{O}_{\mathbb{P}^N}(t + 1 - m - r)) - \dim H^0(\mathbb{P}^N, \text{Im } \alpha_2(t)), \\ & \quad \vdots \\ & \dim H^0(\mathbb{P}^N, \text{Im } \alpha_{d-1}(t)) = b_{d-1} \cdot \dim H^0(\mathbb{P}^N, \mathcal{O}_{\mathbb{P}^N}(t + (d - 1) \cdot (1 - m) - r)) \\ & \quad - b_d \cdot \dim H^0(\mathbb{P}^N, \mathcal{O}_{\mathbb{P}^N}(t + d \cdot (1 - m) - r)). \end{aligned}$$

This shows (b). For $d = N$ we already know that

$$\begin{aligned} \Omega_{\mathbb{P}^N}^T(\log)(t) &\cong \Omega_{\mathbb{P}^N}^N(\log D) \otimes \dots \otimes \Omega_{\mathbb{P}^N}^N(\log D) \otimes \Omega_{\mathbb{P}^N}^{T'}(\log D)(t) \\ &\cong \Omega_{\mathbb{P}^N}^{T'}(\log D)(t + l_N \cdot (m - N - 1)), \end{aligned}$$

which proves assertion (c). In order to prove (d), we consider again the short exact sequences of (3.7) and obtain

$$\begin{aligned} H^q(\mathbb{P}^N, \Omega_{\mathbb{P}^N}^T(\log D)(t)) &= 0 \quad \text{if} \\ H^q(\mathbb{P}^N, \mathcal{O}_{\mathbb{P}^N}(t - r)) &= 0 \quad \text{and} \quad H^{q+1}(\mathbb{P}^N, \text{Im } \alpha_1(t)) = 0, \\ H^{q+1}(\mathbb{P}^N, \text{Im } \alpha_1(t)) &= 0 \quad \text{if} \\ H^{q+1}(\mathbb{P}^N, \mathcal{O}_{\mathbb{P}^N}(t + 1 - m - r)) &= 0 \quad \text{and} \quad H^{q+2}(\mathbb{P}^N, \text{Im } \alpha_2(t)) = 0, \\ &\vdots \end{aligned}$$

$$\begin{aligned} H^{q+d-1}(\mathbb{P}^N, \text{Im } \alpha_{d-1}(t)) &= 0 \quad \text{if} \\ H^{q+d-1}(\mathbb{P}^N, \mathcal{O}_{\mathbb{P}^N}(t + (d - 1) \cdot (1 - m) - r)) &= 0, \end{aligned}$$

and $H^{q+d}(\mathbb{P}^N, \mathcal{O}_{\mathbb{P}^N}(t + d \cdot (1 - m) - r)) = 0$.

This implies $H^q(\mathbb{P}^N, \Omega_{\mathbb{P}^N}^T(\log D)(t)) = 0$ for $1 \leq q \leq N - d - 1$ and $\forall t \in \mathbb{Z}$.

The last statement can be proven by Serre duality which means

$$\begin{aligned} \dim H^q(\mathbb{P}^N, \Omega_{\mathbb{P}^N}^T(\log D)(t)) \\ = \dim H^{N-q}(\mathbb{P}^N, \Omega_{\mathbb{P}^N}^{T^*}(\log D)(-t - m - (l - 1) \cdot (m - N - 1))), \end{aligned}$$

where $\text{depth } T^* = N - d_l < N$. Note if we use (b) with T^* instead of T , we obtain a formula for $\dim H^N(\mathbb{P}^N, \Omega_{\mathbb{P}^N}^T(\log D)(t))$. □

3.3. Symmetric Differential Forms

Let T be a Young tableau with r boxes and only one row, i.e. $\text{depth } T = 1$. We will specify the dimensions of $H^q(\mathbb{P}^N, S^r \Omega^1(\log D)(t))$ and consider the following exact sequence (cf. Lemma 3.5)

$$0 \longrightarrow \bigoplus_{b_1} \mathcal{O}_{\mathbb{P}^N}(-m + 1 - r) \longrightarrow \bigoplus_{b_0} \mathcal{O}_{\mathbb{P}^N}(-r) \longrightarrow S^r \Omega_{\mathbb{P}^N}^1(\log D) \longrightarrow 0 \quad (3.9)$$

with the integers $b_0 = \binom{N+r}{N}$ and $b_1 = \binom{N+r-1}{N}$.

Theorem 3.7. *Let $N \geq 2$. Then one has:*

$$(a) \chi(\mathbb{P}^N, S^r \Omega_{\mathbb{P}^N}^1(\log D)(t)) = \frac{1}{N!} \cdot \binom{N+r}{N} \cdot \prod_{j=1}^N (t-r+j) - \frac{1}{N!} \cdot \binom{N+r-1}{N}$$

$$\begin{aligned}
& \cdot \prod_{i=1}^N (t - m + 1 - r + i). \\
& (b) \dim H^0(\mathbb{P}^N, S^r \Omega_{\mathbb{P}^N}^1(\log D)(t)) = \binom{N+r}{N} \cdot \binom{t-r+N}{N} - \binom{N+r-1}{N} \cdot \binom{t-m+1-r+N}{N}. \\
& (c) \text{ For } 1 \leq q \leq N-2 \text{ it holds: } H^q(\mathbb{P}^N, S^r \Omega_{\mathbb{P}^N}^1(\log D)(t)) = 0 \text{ for all } t \in \mathbb{Z}. \\
& (d) \dim H^{N-1}(\mathbb{P}^N, S^r \Omega_{\mathbb{P}^N}^1(\log D)(t)) \\
& = \sum_{i=0}^{N-1} (-1)^i \cdot \tilde{b}_i \cdot \binom{-t - r(m-2) - i \cdot (m-1) - 1}{N} \\
& - \binom{N+r}{N} \cdot \binom{-t+r-1}{N} + \binom{N+r-1}{N} \cdot \binom{-t+m+r-2}{N} \\
& \dim H^N(\mathbb{P}^N, S^r \Omega_{\mathbb{P}^N}^1(\log D)(t)) \\
& = \sum_{i=0}^{N-1} (-1)^i \cdot \tilde{b}_i \cdot \binom{-t - r(m-2) - i \cdot (m-1) - 1}{N} \text{ with the integers} \\
& \tilde{b}_i = \frac{1}{N+r} \cdot \binom{N+r}{N-1-i} \cdot \binom{N+r}{N} \cdot \binom{r+i-1}{i}. \tag{3.10}
\end{aligned}$$

Proof. (a) follows directly from (3.9) and the additivity of the Euler characteristic. We consider (3.9) together with the corresponding cohomology sequence and know that $H^q(\mathbb{P}^N, \mathcal{O}_{\mathbb{P}^N}(t)) = 0$ for any $q \in \{1, \dots, N-1\}$ and for all $t \in \mathbb{Z}$, which implies (b) and (c). Using the Serre Duality yields

$$\begin{aligned}
& \dim H^N(\mathbb{P}^N, S^r \Omega^1(\log D)(t)) \\
& = \dim H^0(\mathbb{P}^N, \Omega^{T^*}(\log D)(-t + (r-1) \cdot (N+1) - r \cdot m)),
\end{aligned}$$

where T^* is a rectangle with depth $T^* = N-1$ rows and length $T^* = r$ columns and with the associated integers \tilde{b}_i in (3.10) (cf. Lemma 3.5). Theorem 3.6(b) delivers the formula for $\dim H^0(\mathbb{P}^N, \Omega^{T^*}(\log D)(-t + (r-1) \cdot (N+1) - r \cdot m))$.

Finally, one gets easily the dimension $\dim H^{N-1}(\mathbb{P}^N, S^r \Omega_{\mathbb{P}^N}^1(\log D)(t))$ from the long cohomology sequence. \square

Corollary 3.8. *For $N \geq 2$ we obtain:*

- (a) $H^0(\mathbb{P}^N, S^r \Omega_{\mathbb{P}^N}^1(\log D)(t)) \neq 0 \Leftrightarrow t \geq r$.
- (b) $\dim H^0(\mathbb{P}^N, S^r \Omega_{\mathbb{P}^N}^1(\log D)(t)) = \chi(\mathbb{P}^N, S^r \Omega_{\mathbb{P}^N}^1(\log D)(t))$ if $t \geq m+r-N-1$.
- (c) $H^N(\mathbb{P}^N, S^r \Omega_{\mathbb{P}^N}^1(\log D)(t)) = 0 \Leftrightarrow t \geq -r(m-2) - N$.
- (d) $H^{N-1}(\mathbb{P}^N, S^r \Omega_{\mathbb{P}^N}^1(\log D)(t)) = 0$ if $t \geq m+r-N-1$.

Proof. Obviously, the proof follows from Theorem 3.7. \square

Theorem 3.9. *Let $N \geq 2$ and let D be a hyperplane, that is, $m = 1$.*

Then one has

$$(a) \dim H^0(\mathbb{P}^N, S^r \Omega_{\mathbb{P}^N}^1(\log D)(t)) = \binom{N+r-1}{N-1} \cdot \binom{t-r+N}{N}.$$

$$(b) \text{ For } 1 \leq q \leq N-1 \text{ it holds: } H^q(\mathbb{P}^N, S^r \Omega_{\mathbb{P}^N}^1(\log D)(t)) = 0 \quad \forall t \in \mathbb{Z}.$$

$$(c) \dim H^N(\mathbb{P}^N, S^r \Omega_{\mathbb{P}^N}^1(\log D)(t)) = \binom{N+r-1}{N-1} \cdot \binom{-t+r-1}{N}$$

Proof. $S^r \Omega_{\mathbb{P}^N}^1(\log D) \cong \bigoplus_{\binom{N+r-1}{N-1}} \mathcal{O}_{\mathbb{P}^N}(-r).$ □

4. Complete Intersections $Y \subset \mathbb{P}^N$

Let $Y = H_1 \cap \dots \cap H_{N-n} \subseteq \mathbb{P}^N$ be a nonsingular, irreducible, complete intersection of algebraic hypersurfaces $H_i \subset \mathbb{P}^N$, where H_i is given by the equation $F_i = 0$ with $\deg F_i = m_i$. We denote by n the dimension of Y . Let D be a prime divisor on Y , which is defined by the equation $D = Y \cap H$ with a hypersurface $H : F = 0$. The degree of H is m . In the following, we abbreviating denote $c = N - n = \text{codim } Y$ and assume $n \geq 2$. Let X be a further complete intersection which is described by $X = H_1 \cap \dots \cap H_{c-1}$. Here $\dim X = n + 1$ and $Y = X \cap H_c$. There exists also a divisor $D^* = X \cap H$ on X . Assume that the hypersurfaces H_1, \dots, H_{N-n} and H lie in general position, i.e. for instance $X = H_1 \cap \dots \cap H_{c-1} \subseteq \mathbb{P}^N$ and the prime divisors D on Y and D^* on X are nonsingular, irreducible, complete intersections, too.

4.1. Alternating Differential Forms

In case $r = n$ we obtain $\Omega_Y^n = \omega_Y \cong \mathcal{O}_Y(\sum_{i=1}^c m_i - N - 1)$ which implies

$$\Omega_Y^n(\log D) \cong \Omega_Y^n(m) \cong \mathcal{O}_Y\left(\sum_{i=1}^c m_i - N - 1 + m\right),$$

where $D = Y \cap H$ with $\deg H = m$. The dimensions of $H^q(Y, \Omega_Y^n(\log D)(t)) = H^q(Y, \mathcal{O}_Y(\sum_{i=1}^c m_i - N - 1 + m + t))$ are well known:

If $1 \leq q \leq n - 1$ then $H^q(Y, \mathcal{O}_Y(t)) = 0 \quad \forall t \in \mathbb{Z}$.

$$\dim H^0(Y, \mathcal{O}_Y(t)) = \binom{t+N}{N}$$

$$+ \sum_{j=1}^c (-1)^j \cdot \sum_{1 \leq i_1 < i_2 < \dots < i_j \leq c} \binom{t + N - m_{i_1} - m_{i_2} - \dots - m_{i_j}}{N}$$

$$\dim H^n(Y, \mathcal{O}_Y(t)) = \dim H^0(Y, \mathcal{O}_Y(-t + m_1 + m_2 + \dots + m_c - N - 1)),$$

(cf. e.g. [1] or the proof of Lemma (4.4) in the present paper).

We study the cohomology groups $H^q(Y, \Omega_Y^r(\log D)(t))$ with $r < \dim Y = n$:

Lemma 4.1. *The following sequences are exact.*

$$(a) \quad 0 \rightarrow \mathcal{O}_X(-m_c) \rightarrow \mathcal{O}_X \xrightarrow{\varphi^*} \mathcal{O}_Y \rightarrow 0. \quad (4.1)$$

$$(b) \quad 0 \rightarrow \Omega_X^r(\log D^*)(-m_c) \xrightarrow{\alpha} \Omega_X^r(\log D^*) \xrightarrow{\beta} \mathcal{O}_Y \otimes_{\mathcal{O}_X} \Omega_X^r(\log D^*) \rightarrow 0. \quad (4.2)$$

$$(c) \quad 0 \rightarrow \Omega_Y^{r-1}(\log D)(-m_c) \xrightarrow{\gamma} \mathcal{O}_Y \otimes_{\mathcal{O}_X} \Omega_X^r(\log D^*) \xrightarrow{\delta} \Omega_Y^r(\log D) \rightarrow 0. \quad (4.3)$$

Proof. Notice, for $r = 1$ we have to substitute $\Omega_Y^{r-1}(\log D)$ by the structure sheaf \mathcal{O}_Y . The composition $\delta \circ \beta$ is the restriction of the differential forms on X to the subvariety $Y \subset X$. Obviously, the sequence (4.1) is exact and (4.2) results by multiplication of (4.1) with the locally free sheaf $\Omega_X^r(\log D^*)$. We will show that (4.3) is also an exact sequence. Let $U \subseteq X$ be an open subset of X and let $V = Y \cap U$ be an open, nonempty subset of Y . Without loss of generality we assume $U \subseteq U_i = \{x_i \neq 0\}$. Moreover, we suppose the existence of local parameters $u_1, \dots, u_{n-1}, u_n = \frac{F}{x_i^m}, u_{n+1} = \frac{F}{x_i^{m_c}}$ of X on U such that their restriction to Y are also local parameters $v_1 = \varphi^*(u_1), \dots, v_{n-1} = \varphi^*(u_{n-1}), v_n = \varphi^*(u_n) = \frac{F}{x_i^m}$ of Y on V . Then $\Gamma(V, \mathcal{O}_Y \otimes_{\mathcal{O}_X} \Omega_X^r(\log D^*))$ is a free $\Gamma(V, \mathcal{O}_Y)$ -module whose rank is equal to $\binom{n+1}{r}$. Let $\omega \in \Gamma(V, \mathcal{O}_Y \otimes_{\mathcal{O}_X} \Omega_X^r(\log D^*))$ be a section of the form

$$\begin{aligned} \omega &= \sum_{i_\nu=1}^{n-1} f_{i_1, \dots, i_r} \, d u_{i_1} \wedge \dots \wedge d u_{i_r} \\ &\quad + \sum_{i_\nu=1}^{n-1} f_{i_1, \dots, i_{r-1}, n} \, d u_{i_1} \wedge \dots \wedge d u_{i_{r-1}} \wedge \frac{d u_n}{u_n} \\ &\quad + \sum_{i_\nu=1}^{n-1} f_{i_1, \dots, i_{r-1}, n+1} \, d u_{i_1} \wedge \dots \wedge d u_{i_{r-1}} \wedge d u_{n+1} \\ &\quad + \sum_{i_\nu=1}^{n-1} f_{i_1, \dots, i_{r-2}, n, n+1} \, d u_{i_1} \wedge \dots \wedge d u_{i_{r-2}} \wedge \frac{d u_n}{u_n} \wedge d u_{n+1} \end{aligned}$$

where $f_{i_1, \dots, i_r} \in \Gamma(V, \mathcal{O}_Y)$. The homomorphism δ is defined as follows:

$$\delta(\omega) = \sum_{i_\nu=1}^{n-1} f_{i_1, \dots, i_r} dv_{i_1} \wedge \dots \wedge dv_{i_r} + \sum_{i_\nu=1}^{n-1} f_{i_1, \dots, i_{r-1}, n} dv_{i_1} \wedge \dots \wedge dv_{i_{r-1}} \wedge \frac{dv_n}{v_n},$$

which means that $\delta(\omega) \in \Gamma(V, \Omega_Y^r(\log D))$. The kernel of δ is given by

$$\ker \delta = \left\{ \sum_{i_\nu=1}^{n-1} f_{i_1, \dots, i_{r-1}, n+1} du_{i_1} \wedge \dots \wedge du_{i_{r-1}} \wedge du_{n+1} + \sum_{i_\nu=1}^{n-1} f_{i_1, \dots, i_{r-2}, n, n+1} du_{i_1} \wedge \dots \wedge du_{i_{r-2}} \wedge \frac{du_n}{u_n} \wedge du_{n+1} \right\},$$

where $\ker \delta \subseteq \Gamma(V, \mathcal{O}_Y \otimes_{\mathcal{O}_X} \Omega_X^r(\log D^*))$. In order to show that the kernel of δ is isomorphic to $\Gamma(V, \Omega_Y^{r-1}(\log D)(-m_c))$, we consider the following homomorphisms

$$\ker \delta \xrightarrow{\tilde{\alpha}} \Gamma(V, \mathcal{O}_Y(-m_c) \otimes_{\mathcal{O}_X} \Omega_X^{r-1}(\log D^*)) \xrightarrow{\tilde{\beta}} \Gamma(V, \Omega_Y^{r-1}(\log D)(-m_c)).$$

Let $\xi \in \ker \delta$ be any element. The mappings $\tilde{\alpha}$ and $\tilde{\beta}$ are illustrated by

$$\begin{aligned} \tilde{\alpha}(\xi) &= \frac{1}{x_i^{m_c}} \sum_{i_\nu=1}^{n-1} f_{i_1, \dots, i_{r-1}, n+1} du_{i_1} \wedge \dots \wedge du_{i_{r-1}} \\ &+ \frac{1}{x_i^{m_c}} \sum_{i_\nu=1}^{n-1} f_{i_1, \dots, i_{r-2}, n, n+1} du_{i_1} \wedge \dots \wedge du_{i_{r-2}} \wedge \frac{du_n}{u_n}, \end{aligned}$$

respectively,

$$\begin{aligned} \tilde{\beta}(\tilde{\alpha}(\xi)) &= \frac{1}{x_i^{m_c}} \sum_{i_\nu=1}^{n-1} f_{i_1, \dots, i_{r-1}, n+1} dv_{i_1} \wedge \dots \wedge dv_{i_{r-1}} \\ &+ \frac{1}{x_i^{m_c}} \sum_{i_\nu=1}^{n-1} f_{i_1, \dots, i_{r-2}, n, n+1} dv_{i_1} \wedge \dots \wedge dv_{i_{r-2}} \wedge \frac{dv_n}{v_n}. \end{aligned}$$

Since $x_i^{m_c} \cdot du_{n+1} = x_i^{m_c} \cdot d \frac{F_c}{x_i^{m_c}}$ is a global section of the sheaf $\mathcal{O}_Y(m_c) \otimes_{\mathcal{O}_X} \Omega_X^1$ the functions $\tilde{\alpha}$ and $\tilde{\beta}$ are independent of the index i with $U \subseteq U_i$ and independent of the choice of the local parameters u_1, \dots, u_{n-1} . One can easily see that $\tilde{\alpha}$ and $\tilde{\beta}$ are monomorphic. The mapping $\tilde{\beta}$ is the restriction from X to Y which obviously is epimorphic. While $\tilde{\alpha}$ is generally not epimorphic, any element of $\Gamma(V, \Omega_Y^{r-1}(\log D)(-m_c))$ has a preimage in $\ker \delta$. We can represent an element of $\Gamma(V, \Omega_Y^{r-1}(\log D)(-m_c))$ by the form $\tilde{\beta}(\tilde{\alpha}(\xi))$ with functions $f_{i_1, \dots, i_r} \in \Gamma(V, \mathcal{O}_Y)$. In order to find a preimage in $\ker \delta$, we use the same functions f_{i_1, \dots, i_r} , and in place of v_i we take the local parameters u_i on X and multiply

with $x_i^{m_c} \cdot du_{n+1}$. This proves that the composition $\tilde{\beta} \circ \tilde{\alpha}$ is isomorphic, the sequence (4.3) is exact. \square

By means of these exact sequences we are going to prove recursion formulas about the dimensions of the cohomology groups $H^q(Y, \Omega_Y^r(\log D)(t))$. As above mentioned, for $r = n$ these dimensions are known.

Theorem 4.2. (a) $\chi(Y, \Omega_Y^r(\log D)(t)) = \chi(X, \Omega_X^r(\log D^*)(t)) - \chi(X, \Omega_X^r(\log D^*)(t - m_c)) - \chi(Y, \Omega_Y^{r-1}(\log D)(t - m_c))$ for $r \geq 1$

In the case $r = 1$ one has to substitute $\Omega_Y^{r-1}(\log D)$ by the structure sheaf \mathcal{O}_Y .

(b) Let $0 < q < n$, $q+r \neq n$ and $r \geq 0$. Then one has $H^q(Y, \Omega_Y^r(\log D)(t)) = 0$ for any $t \in \mathbb{Z}$.

(c) $\dim H^0(Y, \Omega_Y^r(\log D)(t)) = \dim H^0(X, \Omega_X^r(\log D^*)(t)) - \dim H^0(X, \Omega_X^r(\log D^*)(t - m_c)) - \dim H^0(Y, \Omega_Y^{r-1}(\log D)(t - m_c))$ for $0 < r < n$.

(d) $\dim H^n(Y, \Omega_Y^r(\log D)(t)) = \dim H^0(X, \Omega_X^{n-r}(\log D^*)(-t - m)) - \dim H^0(X, \Omega_X^{n-r}(\log D^*)(-t - m_c - m)) - \dim H^0(Y, \Omega_Y^{n-r-1}(\log D)(-t - m_c - m))$.

(e) $\dim H^1(Y, \Omega_Y^{n-1}(\log D)(t)) = \dim H^0(Y, \Omega_Y^{n-1}(\log D)(t)) + \dim H^0(Y, \Omega_Y^n(\log D)(t + m_c)) + \dim H^0(X, \Omega_X^n(\log D^*)(t)) - \dim H^0(X, \Omega_X^n(\log D^*)(t + m_c)) - \dim H^1(X, \Omega_X^n(\log D^*)(t)) + \dim H^1(X, \Omega_X^n(\log D^*)(t + m_c))$.

(f) $\dim H^{n-r}(Y, \Omega_Y^r(\log D)(t)) = \dim H^{n-r-1}(Y, \Omega_Y^{r+1}(\log D)(t + m_c)) - \dim H^{n-r}(X, \Omega_X^{r+1}(\log D^*)(t)) + \dim H^{n-r}(X, \Omega_X^{r+1}(\log D^*)(t + m_c))$ for $2 \leq r < n$.

Proof. Under the additional condition $q + r < n$ the proof of (b) will be shown by complete induction with respect to $c = \text{codim } Y$ and r . Then the case $q + r > n$ follows directly from the Serre duality. If $c = 0$, i.e. $Y = \mathbb{P}^N$, Theorem 3.3 implies $H^q(Y, \Omega_Y^r(\log D)(t)) = 0$ for $0 < q < N$ and $q + r \neq N$.

If $r = 0$ then we get $H^q(Y, \mathcal{O}_Y(t)) = 0$ for $0 < q < n$ (cf. e.g. [2, Lemma 1]).

In particular, we have the following induction assumption ($c-1 = \text{codim } X$):

(i) From $q, r \in \mathbb{N}$, $0 < q$, $0 \leq r$ and $q + r < n + 1$ it follows

$$H^q(X, \Omega_X^r(\log D^*)(t)) = 0 \text{ for all } t \in \mathbb{Z}.$$

Now assume $0 < q$, $0 \leq r$ and $q + r < n$. From (4.2) we get the exact sequence

$$\dots \longrightarrow H^q(X, \Omega_X^r(\log D^*)(t)) \longrightarrow H^q(Y, \mathcal{O}_Y(t) \otimes_{\mathcal{O}_X} \Omega_X^r(\log D^*)) \longrightarrow$$

$$\longrightarrow H^{q+1}(X, \Omega_X^r(\log D^*)(t - m_c)) \longrightarrow \dots$$

Since $0 < q + 1 + r < n + 1$ we have by induction assumption (i):

$$H^q(X, \Omega_X^r(\log D^*)(t)) = 0 \quad \text{and} \quad H^{q+1}(X, \Omega_X^r(\log D^*)(t - m_c)) = 0.$$

Hence, $H^q(Y, \mathcal{O}_Y(t) \otimes_{\mathcal{O}_X} \Omega_X^r(\log D^*)) = 0$ for $0 < q + r < n$ and any $t \in \mathbb{Z}$.

Now, let $r > 0$ be a fixed integer. We use the following induction assumption:

(ii) If $0 < q$ and $q + r - 1 < n$ then $H^q(Y, \Omega_Y^{r-1}(\log D)(t)) = 0$ for all $t \in \mathbb{Z}$.

To prove: If $0 < q$ and $q + r < n$ then $H^q(Y, \Omega_Y^r(\log D)(t)) = 0$ for all $t \in \mathbb{Z}$. Let $0 < q + r < n$. We consider the exact sequence which is given by (4.3)

$$\begin{aligned} \dots \longrightarrow H^q(Y, \mathcal{O}_Y(t) \otimes_{\mathcal{O}_X} \Omega_X^r(\log D^*)) &\longrightarrow H^q(Y, \Omega_Y^r(\log D)(t)) \longrightarrow \\ &\longrightarrow H^{q+1}(Y, \Omega_Y^{r-1}(\log D)(t - m_c)) \longrightarrow \dots \end{aligned}$$

By (ii) one has $H^{q+1}(Y, \Omega_Y^{r-1}(\log D)(t - m_c)) = 0$ for all $t \in \mathbb{Z}$ since $q + 1 + r - 1 = q + r < n$ (and $q + 1 < n$). Furthermore, we know that $H^q(Y, \mathcal{O}_Y(t) \otimes_{\mathcal{O}_X} \Omega_X^r(\log D^*)) = 0$ for any $t \in \mathbb{Z}$ because of $0 < q + r < n$.

This implies $H^q(Y, \Omega_Y^r(\log D)(t)) = 0$ for $0 < q < n$ and $q + r < n$ for any $t \in \mathbb{Z}$.

For the proof of (c) we first consider the exact sequence from (4.2)

$$\begin{aligned} 0 \longrightarrow H^0(X, \Omega_X^r(\log D^*)(t - m_c)) &\longrightarrow H^0(X, \Omega_X^r(\log D^*)(t)) \longrightarrow \\ \longrightarrow H^0(Y, \mathcal{O}_Y(t) \otimes_{\mathcal{O}_X} \Omega_X^r(\log D^*)) &\longrightarrow H^1(X, \Omega_X^r(\log D^*)(t - m_c)) \longrightarrow \dots, \end{aligned} \tag{4.4}$$

and apply (i) which yields $H^1(X, \Omega_X^r(\log D^*)(t - m_c)) = 0$ as $1 + r < n + 1 = \dim X$. Because of (4.3) one gets the exact sequence

$$\begin{aligned} 0 \longrightarrow H^0(Y, \Omega_Y^{r-1}(\log D)(t - m_c)) &\longrightarrow H^0(Y, \mathcal{O}_Y(t) \otimes_{\mathcal{O}_X} \Omega_X^r(\log D^*)) \longrightarrow \\ H^0(Y, \Omega_Y^r(\log D)(t)) &\longrightarrow H^1(Y, \Omega_Y^{r-1}(\log D)(t - m_c)) \longrightarrow \dots, \end{aligned} \tag{4.5}$$

and due to $1 + r - 1 = r < n$ one has $H^1(Y, \Omega_Y^{r-1}(\log D)(t - m_c)) = 0$. Statement (c) can be read from (4.4) and (4.5). Assertion (d) can easily be shown by Serre duality. The Euler-Poincare characteristic can be calculated by the exact sequences (4.1)–(4.3). This allows us to specify finally the dimension of $H^{n-r}(Y, \Omega_Y^r(\log D)(t))$. (e) and (f) also can be shown using the exact cohomology sequences. \square

4.2. T -Symmetric Differential Forms

Let $Y \subseteq \mathbb{P}^N$ be the n -dimensional complete intersection of multidegree $(m) = (m_1, m_2, \dots, m_c)$. $c = N - n$ denotes the codimension of Y .

We consider a Young tableau T with r boxes, the row lengths $l_1 \geq l_2 \geq \dots \geq l_d > 0$ and the column lengths $d_1 \geq d_2 \geq \dots \geq d_l > 0$. We denote $l = l_1 = \text{length } T$ and $d = d_1 = \text{depth } T$. Let $M(T)$ be the set of all integer matrices $A = ((d_{i,j})) \in \mathbb{N}^{(c+1, l)}$ with $c+1$ rows, l columns and with the following properties:

- (1) $d_{1,j} = d_j \quad \forall j \in \{1, \dots, l\}$,
- (2) $d_{i,l} \geq d_{i+1,l} \geq 0 \quad \forall i \in \{1, \dots, c\}$,
- (3) $d_{i,j} \geq d_{i+1,j} \geq d_{i,j+1} \quad \forall i \in \{1, \dots, c\} \quad \forall j \in \{1, \dots, l-1\}$.

Let $\varrho_i(A) = \sum_{j=1}^l d_{ij}$ be the i -th row sum of A and we put $\varrho(A) = \varrho_{c+1}(A)$. We denote by

$$\mu = \sum_{j=1}^c d_j \tag{4.6}$$

the number of boxes in the first c columns of T , where $d_j = 0$ for $j > l$. One can easily see that $r - \mu \leq \varrho(A) \leq r$ for all $A \in M(T)$. Finally, we define the subset $M_s(T)$ of $M(T)$ by $M_s(T) := \{A \in M(T) : \varrho(A) = r - s\}$ for all $s \in \{0, 1, \dots, \mu\}$. For simplification we set furthermore:

$$\Omega_{\mathbb{P}^N|_Y}^{T'}(\log D^*) = \mathcal{O}_Y \otimes_{\mathcal{O}_{\mathbb{P}^N}} \Omega_{\mathbb{P}^N}^{T'}(\log D^*), \quad E_T^s = \bigoplus_{A \in M_s(T)} \Omega_{\mathbb{P}^N|_Y}^{T'(A)}(\log D^*)(t(A))$$

with $t(A) = \sum_{i=1}^c (\varrho_{i+1}(A) - \varrho_i(A)) \cdot m_i$. Here $T'(A)$ denotes a Young tableau with $\varrho(A)$ boxes and the column lengths $d_{c+1,1}, \dots, d_{c+1,l}$, that is, $T'(A)$ depends only on the last row of A . If $\varrho(A) = 0$ we need to replace the sheaf $\Omega_{\mathbb{P}^N|_Y}^{T'(A)}(\log D^*)$ by the structure sheaf \mathcal{O}_Y .

Lemma 4.3. *There exists following exact sequence:*

$$0 \longrightarrow E_T^\mu \xrightarrow{\beta_\mu} E_T^{\mu-1} \xrightarrow{\beta_{\mu-1}} \dots \xrightarrow{\beta_2} E_T^1 \xrightarrow{\beta_1} \Omega_{\mathbb{P}^N|_Y}^T(\log D^*) \xrightarrow{\beta_0} \Omega_Y^T(\log D) \longrightarrow 0. \tag{4.7}$$

Proof. (4.7) is the T -Power of the following short exact sequence (cf. [3]):

$$0 \longrightarrow \bigoplus_{i=1}^c \mathcal{O}_Y(-m_i) \xrightarrow{\alpha} \mathcal{O}_Y \otimes_{\mathcal{O}_{\mathbb{P}^N}} \Omega_{\mathbb{P}^N}^1(\log D^*) \xrightarrow{\beta} \Omega_Y^1(\log D) \longrightarrow 0. \tag{4.8}$$

We need to show that (4.8) is an exact sequence. Let $U \subseteq \mathbb{P}^N$ be an open subset. Without loss of generality we put $U \subseteq U_i = \{x_i \neq 0\}$. Assume that there exist local parameters $\frac{F_1}{x_i^{m_1}}, \dots, \frac{F_c}{x_i^{m_c}}, u_1, \dots, u_{n-1}, \frac{F}{x_i^m}$ of \mathbb{P}^N on U such that the restrictions $v_1 = \varphi^*(u_1), \dots, v_{n-1} = \varphi^*(u_{n-1}), v_n = \varphi^*\left(\frac{F}{x_i^m}\right)$ are local parameters of Y on $U \cap Y$. We know that $\Gamma(U \cap Y, \mathcal{O}_Y \otimes_{\mathcal{O}_{\mathbb{P}^N}} \Omega_{\mathbb{P}^N}^1(\log D^*))$ is a free $\Gamma(U \cap Y, \mathcal{O}_Y)$ -module defined by the span

$$d \frac{F_1}{x_i^{m_1}}, \dots, d \frac{F_c}{x_i^{m_c}}, d u_1, \dots, d u_{n-1}, \frac{x_i^m}{F} \cdot d \frac{F}{x_i^m} .$$

$\Gamma(U \cap Y, \Omega_Y^1(\log D))$ is a free $\Gamma(U \cap Y, \mathcal{O}_Y)$ -module with the span $d v_1, \dots, d v_n$. Let $\omega \in \Gamma(U \cap Y, \mathcal{O}_Y \otimes_{\mathcal{O}_{\mathbb{P}^N}} \Omega_{\mathbb{P}^N}^1(\log D^*))$ be any element given by

$$\omega = \sum_{j=1}^c f_j \cdot x_i^{m_j} \cdot d \frac{F_j}{x_i^{m_j}} + \sum_{k=1}^{n-1} g_k \cdot d u_k + h \cdot \frac{x_i^m}{F} \cdot d \frac{F}{x_i^m} .$$

The homomorphism β maps ω to $\beta(\omega) = \sum_{k=1}^{n-1} g_k \cdot d v_k + h \cdot \frac{d v_n}{v_n}$, where the kernel of this mapping is given by $\ker \beta = \left\{ \sum_{j=1}^c f_j \cdot x_i^{m_j} \cdot d \frac{F_j}{x_i^{m_j}} \right\}$. We obtain the following homomorphism

$$\gamma : \bigoplus_{j=1}^c \Gamma(U \cap Y, \mathcal{O}_Y(-m_j)) \longrightarrow \ker \beta \quad \text{with} \quad (f_1, \dots, f_c) \longmapsto \sum_{j=1}^c f_j \cdot x_i^{m_j} \cdot d \frac{F_j}{x_i^{m_j}}$$

which is isomorphic and independent of the index i with $U \subseteq U_i$. □

Lemma 4.4. *For an arbitrary Young tableau T' there exists the following exact sequence*

$$\begin{aligned} 0 &\longrightarrow \Omega_{\mathbb{P}^N}^{T'}(\log D^*)\left(-\sum_{i=1}^c m_i\right) \xrightarrow{\alpha_c} \bigoplus_{1 \leq i \leq c} \Omega_{\mathbb{P}^N}^{T'}(\log D^*)\left(-\sum_{j=1}^c m_j + m_i\right) \xrightarrow{\alpha_{c-1}} \dots \\ &\dots \xrightarrow{\alpha_3} \bigoplus_{1 \leq i_1 < i_2 \leq c} \Omega_{\mathbb{P}^N}^{T'}(\log D^*)\left(-m_{i_1} - m_{i_2}\right) \xrightarrow{\alpha_2} \bigoplus_{1 \leq i \leq c} \Omega_{\mathbb{P}^N}^{T'}(\log D^*)\left(-m_i\right) \xrightarrow{\alpha_1} \\ &\xrightarrow{\alpha_1} \Omega_{\mathbb{P}^N}^{T'}(\log D^*) \xrightarrow{\alpha_0} \mathcal{O}_Y \otimes_{\mathcal{O}_{\mathbb{P}^N}} \Omega_{\mathbb{P}^N}^{T'}(\log D^*) \longrightarrow 0. \end{aligned} \tag{4.9}$$

Proof. We consider the following exact sequence which is called the Koszul complex:

$$0 \longrightarrow \mathcal{O}_{\mathbb{P}^N}\left(-\sum_{i=1}^c m_i\right) \xrightarrow{\alpha_c} \bigoplus_{1 \leq i \leq c} \mathcal{O}_{\mathbb{P}^N}\left(-\sum_{j=1}^c m_j + m_i\right) \xrightarrow{\alpha_{c-1}} \dots$$

$$\xrightarrow{\alpha_3} \bigoplus_{1 \leq i_1 < i_2 \leq c} \mathcal{O}_{\mathbb{P}^N}(-m_{i_1} - m_{i_2}) \xrightarrow{\alpha_2} \bigoplus_{1 \leq i \leq c} \mathcal{O}_{\mathbb{P}^N}(-m_i) \xrightarrow{\alpha_1} \mathcal{O}_{\mathbb{P}^N} \xrightarrow{\alpha_0} \mathcal{O}_Y \longrightarrow 0.$$

Multiplying this exact sequence with the local free sheaf $\Omega_{\mathbb{P}^N}^{T'}(\log D^*)$ yields the assertion. \square

Theorem 4.5. *Under the assumption $1 \leq q < n - \text{depth } T - \mu$ one gets $H^q(Y, \Omega_Y^T(\log D)(t)) = 0$ for all $st \in \mathbb{Z}$.*

Proof. We write instead of (4.9) short exact sequences and obtain

$$\begin{aligned} 0 \longrightarrow \Omega_{\mathbb{P}^N}^{T'}(\log D^*)(-\sum_{i=1}^c m_i) &\longrightarrow \bigoplus_{1 \leq i \leq c} \Omega_{\mathbb{P}^N}^{T'}(\log D^*)(-\sum_{j=1}^c m_j + m_i) \longrightarrow \\ &\longrightarrow \text{Im } \alpha_{c-1} \longrightarrow 0, \end{aligned}$$

$$\begin{aligned} 0 \longrightarrow \text{Im } \alpha_{c-1} &\longrightarrow \bigoplus_{1 \leq i_1 < i_2 \leq c} \Omega_{\mathbb{P}^N}^{T'}(\log D^*)(-\sum_{j=1}^c m_j + m_{i_1} + m_{i_2}) \longrightarrow \\ &\longrightarrow \text{Im } \alpha_{c-2} \longrightarrow 0, \end{aligned}$$

$\vdots \qquad \qquad \qquad \vdots$

$$0 \longrightarrow \text{Im } \alpha_2 \longrightarrow \bigoplus_{1 \leq i \leq c} \Omega_{\mathbb{P}^N}^{T'}(\log D^*)(-m_i) \longrightarrow \text{Im } \alpha_1 \longrightarrow 0,$$

$$0 \longrightarrow \text{Im } \alpha_1 \longrightarrow \Omega_{\mathbb{P}^N}^{T'}(\log D^*) \longrightarrow \mathcal{O}_Y \otimes_{\mathcal{O}_{\mathbb{P}^N}} \Omega_{\mathbb{P}^N}^{T'}(\log D^*) \longrightarrow 0.$$

Using the long exact cohomology sequences yields a vanishing criterion for $H^q(Y, \mathcal{O}_Y \otimes_{\mathcal{O}_{\mathbb{P}^N}} \Omega_{\mathbb{P}^N}^{T'}(\log D^*)(t))$. We have

$$\begin{aligned} H^q(Y, \mathcal{O}_Y \otimes_{\mathcal{O}_{\mathbb{P}^N}} \Omega_{\mathbb{P}^N}^{T'}(\log D^*)(t)) = 0 &\text{ if } H^q(\mathbb{P}^N, \Omega_{\mathbb{P}^N}^{T'}(\log D^*)(t)) = 0, \\ &\text{ and } H^{q+1}(\mathbb{P}^N, \text{Im } \alpha_1(t)) = 0, \end{aligned}$$

$$\begin{aligned} H^{q+1}(\mathbb{P}^N, \text{Im } \alpha_1(t)) = 0 &\text{ if } H^{q+1}(\mathbb{P}^N, \bigoplus_{1 \leq i \leq c} \Omega_{\mathbb{P}^N}^{T'}(\log D^*)(t - m_i)) = 0, \\ &\text{ and } H^{q+2}(\mathbb{P}^N, \text{Im } \alpha_2(t)) = 0, \end{aligned}$$

\vdots

$$H^{q+c-1}(\mathbb{P}^N, \text{Im } \alpha_{c-1}(t)) = 0$$

$$\text{if } H^{q+c-1}(\mathbb{P}^N, \bigoplus_{1 \leq i \leq c} \Omega_{\mathbb{P}^N}^{T'}(\log D^*)(t - \sum_{j=1}^c m_j + m_i)) = 0$$

$$\text{and } H^{q+c}(\mathbb{P}^N, \Omega_{\mathbb{P}^N}^{T'}(\log D^*)(t - \sum_{i=1}^c m_i)) = 0.$$

Applying Theorem 3.6 (d) yields $H^q(Y, \mathcal{O}_Y(t) \otimes_{\mathcal{O}_{\mathbb{P}^N}} \Omega_{\mathbb{P}^N}^{T'}(\log D^*)) = 0$ for $1 \leq q < n - \text{depth } T'$. Now we study $H^q(Y, \Omega_Y^T(\log D)(t))$ with the aid of (4.7). Decomposing (4.7) in short exact sequences delivers

$$\begin{aligned} 0 &\longrightarrow E_T^\mu \longrightarrow E_T^{\mu-1} \longrightarrow \text{Im } \beta_{\mu-1} \longrightarrow 0, \\ 0 &\longrightarrow \text{Im } \beta_{\mu-1} \longrightarrow E_T^{\mu-2} \longrightarrow \text{Im } \beta_{\mu-2} \longrightarrow 0, \\ &\qquad \qquad \qquad \vdots \qquad \qquad \qquad \vdots \\ 0 &\longrightarrow \text{Im } \beta_2 \longrightarrow E_T^1 \longrightarrow \text{Im } \beta_1 \longrightarrow 0, \\ 0 &\longrightarrow \text{Im } \beta_1 \longrightarrow \Omega_{\mathbb{P}^N|Y}^T(\log D^*) \longrightarrow \Omega_Y^T(\log D) \longrightarrow 0. \end{aligned}$$

With $E_T^s(t) = \bigoplus_{A \in M_s(T)} \Omega_{\mathbb{P}^N|Y}^{T'(A)}(\log D^*)(t + t(A))$ one has

$$\begin{aligned} H^q(Y, \Omega_Y^T(\log D)(t)) = 0 \text{ if } & H^q(Y, \Omega_{\mathbb{P}^N|Y}^T(\log D^*)(t)) = 0, \\ & \text{and } H^{q+1}(Y, \text{Im } \beta_1(t)) = 0, \\ & \qquad \qquad \qquad \vdots \\ H^{q+\mu-1}(Y, \text{Im } \beta_{\mu-1}(t)) = 0 \text{ if } & H^{q+\mu-1}(Y, E_T^{\mu-1}(t)) = 0, \\ & \text{and } H^{q+\mu}(Y, E_T^\mu(t)) = 0. \end{aligned}$$

This implies $H^q(Y, \Omega_Y^T(\log D)(t)) = 0$ for $1 \leq q < n - \text{depth } T - \mu$. □

Now assume for instance $\mu < n - \text{depth } T$. Then for each $t \in \mathbb{Z}$ it follows from our exact sequences:

$$\begin{aligned} H^q(\mathbb{P}^N, \text{Im } \alpha_i(t)) &= 0 \text{ if } 1 \leq q \leq \mu + i, \\ H^q(Y, \mathcal{O}_Y(t) \otimes_{\mathcal{O}_{\mathbb{P}^N}} \Omega_{\mathbb{P}^N}^{T'}(\log D^*)) &= 0 \text{ if } 1 \leq q \leq \mu + c, \\ H^q(Y, E_T^j(t)) &= 0 \text{ if } 1 \leq q \leq j, \quad H^q(Y, \text{Im } \beta_j(t)) = 0 \text{ if } 1 \leq q \leq j. \end{aligned}$$

In particular, the cohomology groups $H^1(\dots)$ of all these sheaves vanish. Therefore, we have the opportunity to calculate the dimensions of their cohomology groups $H^0(\dots)$: Let $h^T(t)$ abbreviating denotes the dimension $\dim H^0(\mathbb{P}^N, \Omega_{\mathbb{P}^N}^T(\log D^*)(t))$ as an integer function of t . Remember that $(m) = (m_1, m_2, \dots, m_c)$ is the multidegree of the complete intersection Y . We set

$$h_{(m)}^T(t) := h^T(t) + \sum_{s=1}^c (-1)^s \cdot \sum_{1 \leq i_1 < i_2 < \dots < i_s \leq c} h^T(t - m_{i_1} - m_{i_2} - \dots - m_{i_s}).$$

Because of (4.9) we have $\dim H^0(Y, \mathcal{O}_Y \otimes_{\mathcal{O}_{\mathbb{P}^N}} \Omega_{\mathbb{P}^N}^{T'}(\log D^*)(t)) = h_{(m)}^{T'}(t)$ and using (4.7) we get the following formula:

Theorem 4.6. *If $\mu < \dim Y - \text{depth } T$ then $\dim H^0(Y, \Omega_Y^T(\log D)(t)) =$*

$$\sum_{A \in M(T)} (-1)^{r-\varrho(A)} \cdot h_{(m)}^{T'(A)}(t + t(A)) \text{ with } t(A) = \sum_{i=1}^c (\varrho_{i+1}(A) - \varrho_i(A)) \cdot m_i.$$

In particular for $t = 0$: $H^0(Y, \Omega_Y^T(\log D)) = 0$ if $\mu < \dim Y - \text{depth } T$.

Remark 4.7. For regular T -symmetrical tensor differential forms one has $H^0(Y, \Omega_Y^T) = 0$ if $\mu < \dim Y$.

4.3. Symmetric Differential Forms

We consider symmetrical differential forms with logarithmic poles as a special case, that means, T is a Young tableau with r boxes and only one row ($\text{depth } T = 1$ $l = \text{length } T = r$).

Let $D^* = H$ be the prime divisor on projective space \mathbb{P}^N and let D be the prime divisor on the n -dimensional complete intersection Y as above ($n \geq 2$). Distinguishing the cases $r \leq c$ and $c < r$ we obtain two exact sequences as symmetrical power of (4.8): Assume at first $r \leq c$:

$$\begin{aligned} 0 &\longrightarrow \bigoplus_{1 \leq i_1 < i_2 < \dots < i_r \leq c} \mathcal{O}_Y(-m_{i_1} - m_{i_2} - \dots - m_{i_r}) \longrightarrow \\ &\longrightarrow \bigoplus_{1 \leq i_1 < \dots < i_{r-1} \leq c} \mathcal{O}_Y(-m_{i_1} - m_{i_2} - \dots - m_{i_{r-1}}) \otimes_{\mathcal{O}_{\mathbb{P}^N}} \Omega_{\mathbb{P}^N}^1(\log D^*) \longrightarrow \dots \\ &\dots \longrightarrow \bigoplus_{1 \leq i \leq c} \mathcal{O}_Y(-m_i) \otimes_{\mathcal{O}_{\mathbb{P}^N}} S^{r-1} \Omega_{\mathbb{P}^N}^1(\log D^*) \longrightarrow \\ &\longrightarrow \mathcal{O}_Y \otimes_{\mathcal{O}_{\mathbb{P}^N}} S^r \Omega_{\mathbb{P}^N}^1(\log D^*) \longrightarrow S^r \Omega_Y^1(\log D) \longrightarrow 0. \end{aligned}$$

In the case $c < r$ the following sequence is exact:

$$\begin{aligned} 0 &\longrightarrow \mathcal{O}_Y(-\sum_{j=1}^c m_j) \otimes_{\mathcal{O}_{\mathbb{P}^N}} \Omega_{\mathbb{P}^N}^{r-c}(\log D^*) \longrightarrow \\ &\longrightarrow \bigoplus_{1 \leq i \leq c} \mathcal{O}_Y(-\sum_{j=1}^c m_j + m_i) \otimes_{\mathcal{O}_{\mathbb{P}^N}} \Omega_{\mathbb{P}^N}^{r-c+1}(\log D^*) \longrightarrow \dots \\ &\dots \longrightarrow \bigoplus_{1 \leq i \leq c} \mathcal{O}_Y(-m_i) \otimes_{\mathcal{O}_{\mathbb{P}^N}} S^{r-1} \Omega_{\mathbb{P}^N}^1(\log D^*) \longrightarrow \\ &\longrightarrow \mathcal{O}_Y \otimes_{\mathcal{O}_{\mathbb{P}^N}} S^r \Omega_{\mathbb{P}^N}^1(\log D^*) \longrightarrow S^r \Omega_Y^1(\log D) \longrightarrow 0. \end{aligned}$$

Furthermore, we have Lemma 4.4 with the sheaf $S^r \Omega_{\mathbb{P}^N}^1(\log D^*)$ instead of $\Omega_{\mathbb{P}^N}^{T'}(\log D^*)$. With the corresponding cohomology sequences we get:

Theorem 4.8. Assume $n = \dim Y \geq 2$.

(a) If $1 \leq q \leq n - 2$ then $H^q(Y, \mathcal{O}_Y(t) \otimes_{\mathcal{O}_{\mathbb{P}^N}} S^r \Omega_{\mathbb{P}^N}^1(\log D^*)) = 0 \forall t \in \mathbb{Z}$.

(b)

$$\dim H^0(Y, \mathcal{O}_Y(t) \otimes_{\mathcal{O}_{\mathbb{P}^N}} S^r \Omega_{\mathbb{P}^N}^1(\log D^*)) = \dim H^0(\mathbb{P}^N, S^r \Omega_{\mathbb{P}^N}^1(\log D^*)(t)) + \sum_{j=1}^c (-1)^j \cdot \sum_{1 \leq i_1 < \dots < i_j \leq c} \dim H^0(\mathbb{P}^N, S^r \Omega_{\mathbb{P}^N}^1(\log D^*)(t - m_{i_1} - \dots - m_{i_j})).$$

(c) $H^0(Y, \mathcal{O}_Y(t) \otimes_{\mathcal{O}_{\mathbb{P}^N}} S^r \Omega_{\mathbb{P}^N}^1(\log D^*)) \neq 0 \Leftrightarrow t \geq r$.

(d) In case $t = 0$: $H^0(Y, \mathcal{O}_Y \otimes_{\mathcal{O}_{\mathbb{P}^N}} S^r \Omega_{\mathbb{P}^N}^1(\log D^*)) = 0$ for all $r > 0$.

Theorem 4.9.

(a) If $r \leq c$ and $1 \leq q < n - r$ then $H^q(Y, S^r \Omega_Y^1(\log D)(t)) = 0 \forall t \in \mathbb{Z}$.

(b) If $c < r$ and $1 \leq q < n - c - 1$ then $H^q(Y, S^r \Omega_Y^1(\log D)(t)) = 0 \forall t \in \mathbb{Z}$.

Proof. By Theorem 4.5 we know $H^q(Y, \Omega_Y^T(\log D)(t)) = 0$ for all $t \in \mathbb{Z}$ if $1 \leq q < n - \text{depth } T - \mu$. For symmetric differential forms we have $\text{depth } T = 1$ and $\mu = \sum_{i=1}^c d_i = \min\{c, r\}$, where $d_i = 1$ for $i \leq r$ and $d_i = 0$ for $i > r$. This proves (b). Under condition $r \leq c$ one gets the stronger result (a) since $H^q(Y, \mathcal{O}_Y(t)) = 0$ for $1 \leq q < n$ and for all $t \in \mathbb{Z}$. \square

Theorem 4.10.

(c) If $r \leq c$ and $r < n$ then

$$H^0(Y, S^r \Omega_Y^1(\log D)(t)) = \dim H^0(Y, \mathcal{O}_Y(t) \otimes_{\mathcal{O}_{\mathbb{P}^N}} S^r \Omega_{\mathbb{P}^N}^1(\log D^*)) + \sum_{k=1}^{r-1} (-1)^k \cdot \sum_{1 \leq i_1 < \dots < i_k \leq c} \dim H^0(Y, \mathcal{O}_Y(t - \sum_{j=1}^k m_{i_j}) \otimes_{\mathcal{O}_{\mathbb{P}^N}} S^{r-k} \Omega_{\mathbb{P}^N}^1(\log D^*)) + (-1)^r \cdot \sum_{1 \leq i_1 < \dots < i_r \leq c} \dim H^0(Y, \mathcal{O}_Y(t - m_{i_1} - \dots - m_{i_r})).$$

(d) If $c < r$ and $c < n - 1$ then

$$H^0(Y, S^r \Omega_Y^1(\log D)(t)) = \dim H^0(Y, \mathcal{O}_Y(t) \otimes_{\mathcal{O}_{\mathbb{P}^N}} S^r \Omega_{\mathbb{P}^N}^1(\log D^*)) + \sum_{k=1}^c (-1)^k \cdot \sum_{1 \leq i_1 < \dots < i_k \leq c} \dim H^0(Y, \mathcal{O}_Y(t - \sum_{j=1}^k m_{i_j}) \otimes_{\mathcal{O}_{\mathbb{P}^N}} S^{r-k} \Omega_{\mathbb{P}^N}^1(\log D^*)).$$

Proof. Statements (c) and (d) follow from the related exact sequences since under these premises by Theorem 4.8 the cohomology groups $H^1(\dots)$ of all these sheaves vanish (cf. Theorem 4.8 and Theorem 3.7). \square

Finally, it is easy to see:

Theorem 4.11. (e) If $t < r \leq \min(c, n-1)$ then $H^0(Y, S^r \Omega_Y^1(\log D)(t)) = 0$.

(f) If $t < r$ and $c < \min(r, n-1)$ then $H^0(Y, S^r \Omega_Y^1(\log D)(t)) = 0$.

(g) If $c < n-1$ then $H^0(Y, S^r \Omega_Y^1(\log D)) = 0$ for all $r > 0$.

(h) If $0 < r < n$ then $H^0(Y, S^r \Omega_Y^1(\log D)) = 0$.

Remark 4.12. On the other hand, for regular symmetrical differential forms on complete intersections it is well known:

If $c < n$ then $H^0(Y, S^r \Omega_Y^1) = 0$ for all $r > 0$.

References

- [1] P. Brückmann, Zur Kohomologie von projektiven Hyperflächen, *Beiträge zur Algebra und Geometrie*, **2** (1974), 87-101.
- [2] P. Brückmann, Zur Kohomologie von vollständigen Durchschnitten mit Koeffizienten in der Garbe der Keime der Differentialformen, *Mathematische Nachrichten*, **71** (1976), 203-210.
- [3] P. Brückmann, The Hilbert polynomial of the sheaf Ω^T of germs of T -symmetrical tensor differential forms on complete intersections, *Math. Ann.*, **307** (1997), 461-472.
- [4] H. Esnault, E. Viehweg, *Lectures on Vanishing Theorems*, Basel-Boston-Berlin (1992).
- [5] W. Fulton, J. Harris, *Representation Theory*, GTM 129, New York-Berlin-Heidelberg (1991).
- [6] R. Hartshorne, *Algebraic Geometry*, GTM 52, New York-Berlin-Heidelberg (1977).
- [7] J.-P. Serre, Faisceaux algébriques cohérents, *Ann. of Math.*, **61** (1955), 197-278.
- [8] I.R. Shafarevich, *Algebraic Geometry II*, Encyclopaedia of Mathematical Sciences, Vol. 35, Berlin, Heidelberg, New York (1996).