

# EXISTENCE OF SOLUTIONS FOR SINGULAR DOUBLE PHASE PROBLEMS VIA THE NEHARI MANIFOLD METHOD

WULONG LIU, GUOWEI DAI, NIKOLAOS S. PAPAGEORGIU, AND PATRICK WINKERT

ABSTRACT. In this paper we study quasilinear elliptic equations driven by the double phase operator and a right-hand side which has the combined effect of a singular and of a parametric term. Based on the fibering method by using the Nehari manifold we are going to prove the existence of at least two weak solutions for such problems when the parameter is sufficiently small.

## 1. INTRODUCTION

Zhikov [62] was the first who introduced and studied functionals whose integrands change their ellipticity according to a point in order to provide models for strongly anisotropic materials. As a prototype he considered the functional

$$\omega \mapsto \int_{\Omega} (|\nabla\omega|^p + \mu(x)|\nabla\omega|^q) dx, \quad (1.1)$$

where  $1 < p < q$  and with a nonnegative weight function  $\mu \in L^\infty(\Omega)$ . Therefore, the integrand of (1.1) has unbalanced growth. The main feature of the functional defined in (1.1) is the change of ellipticity on the set where the weight function is zero, that is, on the set  $\{x \in \Omega : \mu(x) = 0\}$ . In other words, the energy density of (1.1) exhibits ellipticity in the gradient of order  $q$  on the points  $x$  where  $\mu(x)$  is positive and of order  $p$  on the points  $x$  where  $\mu(x)$  vanishes. We also refer to the book of Zhikov-Kozlov-Oleřnik [63]. Functionals of type (1.1) have been intensively studied in the past decade concerning regularity for isotropic and anisotropic settings. We mention the papers of Baroni-Colombo-Mingione [4, 5, 6], Baroni-Kuusi-Mingione [7], Byun-Oh [9, 10], Byun-Youn [11], Colombo-Mingione [14, 15, 16], De Filippis-Mingione [18, 19, 20], De Filippis-Palatucci [21], Esposito-Leonetti-Mingione [23], Esposito-Leonetti-Petricca [24], Marcellini [37, 38, 39], Ok [44, 45], Ragusa-Tachikawa [54], Riey [55] and the references therein.

The energy functional (1.1) is related to the so-called double phase operator which is defined by

$$\operatorname{div} (|\nabla u|^{p-2}\nabla u + \mu(x)|\nabla u|^{q-2}\nabla u) \quad \text{for } u \in W_0^{1,\mathcal{H}}(\Omega) \quad (1.2)$$

with an appropriate Musielak-Orlicz Sobolev space  $W_0^{1,\mathcal{H}}(\Omega)$ , see its definition in Section 2. It is easy to see that (1.2) reduces to the  $p$ -Laplacian if  $\mu \equiv 0$  or to the weighted  $(q, p)$ -Laplacian if  $\inf_{\overline{\Omega}} \mu \geq \mu_0 > 0$ , respectively.

---

2020 *Mathematics Subject Classification.* 35J15, 35J62, 35J92, 35P30.

*Key words and phrases.* Double phase operator, fibering method, multiple solutions, Nehari manifold, singular problems.

Given a bounded domain  $\Omega \subset \mathbb{R}^N$ ,  $N \geq 2$ , with Lipschitz boundary  $\partial\Omega$ , we study the following singular double phase problem

$$\begin{aligned} -\operatorname{div} (|\nabla u|^{p-2}\nabla u + \mu(x)|\nabla u|^{q-2}\nabla u) &= a(x)u^{-\gamma} + \lambda u^{r-1} && \text{in } \Omega, \\ u &= 0 && \text{on } \partial\Omega, \end{aligned} \quad (1.3)$$

where we suppose the subsequent assumptions:

- (H): (i)  $1 < p < N$ ,  $p < q < p^* = \frac{Np}{N-p}$  and  $0 \leq \mu(\cdot) \in L^\infty(\Omega)$ ;  
(ii)  $0 < \gamma < 1$  and  $q < r < p^*$ ;  
(iii)  $a \in L^\infty(\Omega)$  and  $a(x) > 0$  for a. a.  $x \in \Omega$ .

A function  $u \in W_0^{1,\mathcal{H}}(\Omega)$  is said to be a weak solution if  $a(\cdot)u^{-\gamma}h \in L^1(\Omega)$ ,  $u > 0$  for a. a.  $x \in \Omega$  and

$$\begin{aligned} &\int_{\Omega} (|\nabla u|^{p-2}\nabla u + \mu(x)|\nabla u|^{q-2}\nabla u) \cdot \nabla h \, dx \\ &= \int_{\Omega} a(x)u^{-\gamma}h \, dx + \lambda \int_{\Omega} u^{r-1}h \, dx \end{aligned} \quad (1.4)$$

is satisfied for all  $h \in W_0^{1,\mathcal{H}}(\Omega)$ . Due to (H)(ii) we see that  $\int_{\Omega} u^{r-1}h \, dx$  is finite since  $r < p^*$ . So, the definition of a weak solution in (1.4) is well-defined. The corresponding energy functional  $\varphi_\lambda: W_0^{1,\mathcal{H}}(\Omega) \rightarrow \mathbb{R}$  for problem (1.3) is given by

$$\varphi_\lambda(u) = \frac{1}{p}\|\nabla u\|_p^p + \frac{1}{q}\|\nabla u\|_{q,\mu}^q - \frac{1}{1-\gamma} \int_{\Omega} a(x)|u|^{1-\gamma} \, dx - \frac{\lambda}{r}\|u\|_r^r.$$

The main result in this paper is the following theorem.

**Theorem 1.1.** *Let hypotheses (H) be satisfied. Then there exists  $\hat{\lambda}_0^* > 0$  such that for all  $\lambda \in (0, \hat{\lambda}_0^*]$  problem (1.3) has at least two weak solutions  $u^*, v^* \in W_0^{1,\mathcal{H}}(\Omega)$  such that  $\varphi_\lambda(u^*) < 0 < \varphi_\lambda(v^*)$ .*

The main characteristic in our treatment is the usage of the so-called Nehari manifold which turned into a very powerful tool in order to find solutions for differential equations via critical point theory. This method was first introduced by Nehari [42, 43] and the idea behind is the following: For a real reflexive Banach space  $X$  and a functional  $\Psi \in C^1(X, \mathbb{R})$ , we see that a critical point  $u \neq 0$  of  $\Psi$  belongs to the set

$$\mathcal{N} = \left\{ u \in X \setminus \{0\} : \langle \Psi'(u), u \rangle = 0 \right\},$$

where  $\langle \cdot, \cdot \rangle$  is the duality pairing between  $X$  and its dual space  $X^*$ . Therefore,  $\mathcal{N}$  is a natural constraint for finding nontrivial critical points of  $\Psi$ . We mention the book chapter of Szulkin-Weth [57] in order to have a very well description of the method.

Because of the appearance of the singular term in (1.3), it is clear that the corresponding energy functional for problem (1.3) is not  $C^1$  and so we need to make several modifications in order to use the Nehari manifold method. With our work we extend the recent papers of Papageorgiou-Repovš-Vetro [49] for the weighted  $(p, q)$ -Laplacian and Papageorgiou-Winkert [51] for the  $p$ -Laplacian. In contrast to these works we are working in Musielak-Orlicz Sobolev spaces and not in usual Sobolev spaces.

To the best of our knowledge, there are only two works dealing with singular double phase problems. Chen-Ge-Wen-Cao [12] considered problems of type (1.3) and proved the existence of a weak solution with negative energy. Very recently, Farkas-Winkert [25] studied singular Finsler double phase problems of the form

$$-\operatorname{div}(F^{p-1}(\nabla u)\nabla F(\nabla u) + \mu(x)F^{q-1}(\nabla u)\nabla F(\nabla u)) = u^{p^*-1} + \lambda(u^{\gamma-1} + g(u))$$

in  $\Omega$  and  $u = 0$  on  $\partial\Omega$ , where  $(\mathbb{R}^N, F)$  is a Minkowski space. Based on variational tools, the existence of one weak solution is shown. Both works show only the existence of one weak solution (in contrast to our work) and the treatments are completely different from ours.

Finally, existence results for double phase problems with homogeneous Dirichlet or nonlinear Neumann boundary conditions without singular term can be found in the papers of Colasuonno-Squassina [13], El Manouni-Marino-Winkert [22], Gasiński-Papageorgiou [26], Gasiński-Winkert [28, 29, 30], Liu-Dai [34, 35, 36], Marino-Winkert [40], Papageorgiou-Rădulescu-Repovš [47], Papageorgiou-Vetro-Vetro [50], Perera-Squassina [52], Zeng-Bai-Gasiński-Winkert [60, 61] and the references therein. For related works dealing with certain types of double phase problems we refer to the works of Alves-Santos-Silva [1], Bahrouni-Rădulescu-Winkert [2], Barletta-Tornatore [3], Biagi-Esposito-Vecchi [8], Lei [33], Papageorgiou-Rădulescu-Repovš [46], Rădulescu [53], Sun-Wu-Long [56], Wang-Zhao-Zhao [58] and Zeng-Bai-Gasiński-Winkert [59].

## 2. PRELIMINARIES

In this section we recall the main properties on the theory of Musielak-Orlicz spaces  $L^{\mathcal{H}}(\Omega)$  and  $W_0^{1,\mathcal{H}}(\Omega)$ , respectively. We refer to Colasuonno-Squassina [13], Harjulehto-Hästö [31] and Musielak [41] for the main results in this direction.

We denote by  $L^r(\Omega)$  and  $L^r(\Omega; \mathbb{R}^N)$  the usual Lebesgue spaces equipped with the norm  $\|\cdot\|_r$  for every  $1 \leq r < \infty$ . For  $1 < r < \infty$ ,  $W^{1,r}(\Omega)$  and  $W_0^{1,r}(\Omega)$  stand for the Sobolev spaces endowed with the norms  $\|\cdot\|_{1,r}$  and  $\|\cdot\|_{1,r,0} = \|\nabla \cdot\|_r$ , respectively.

Let  $\mathcal{H}: \Omega \times [0, \infty) \rightarrow [0, \infty)$  be the function defined by

$$\mathcal{H}(x, t) = t^p + \mu(x)t^q.$$

Then, the Musielak-Orlicz space  $L^{\mathcal{H}}(\Omega)$  is defined by

$$L^{\mathcal{H}}(\Omega) = \left\{ u \mid u: \Omega \rightarrow \mathbb{R} \text{ is measurable and } \rho_{\mathcal{H}}(u) < +\infty \right\}$$

equipped with the Luxemburg norm

$$\|u\|_{\mathcal{H}} = \inf \left\{ \tau > 0 : \rho_{\mathcal{H}}\left(\frac{u}{\tau}\right) \leq 1 \right\},$$

where the modular function  $\rho_{\mathcal{H}}: L^{\mathcal{H}}(\Omega) \rightarrow \mathbb{R}$  is given by

$$\rho_{\mathcal{H}}(u) := \int_{\Omega} \mathcal{H}(x, |u|) dx = \int_{\Omega} (|u|^p + \mu(x)|u|^q) dx. \quad (2.1)$$

From Colasuonno-Squassina [13, Proposition 2.14] we know that the space  $L^{\mathcal{H}}(\Omega)$  is a reflexive Banach space. Moreover, we define the seminormed space

$$L_{\mu}^q(\Omega) = \left\{ u \mid u: \Omega \rightarrow \mathbb{R} \text{ is measurable and } \int_{\Omega} \mu(x)|u|^q dx < +\infty \right\},$$

which is endowed with the seminorm

$$\|u\|_{q,\mu} = \left( \int_{\Omega} \mu(x)|u|^q dx \right)^{\frac{1}{q}}.$$

In the same way we define  $L_{\mu}^q(\Omega; \mathbb{R}^N)$ .

The Musielak-Orlicz Sobolev space  $W^{1,\mathcal{H}}(\Omega)$  is defined by

$$W^{1,\mathcal{H}}(\Omega) = \{u \in L^{\mathcal{H}}(\Omega) : |\nabla u| \in L^{\mathcal{H}}(\Omega)\}$$

equipped with the norm

$$\|u\|_{1,\mathcal{H}} = \|\nabla u\|_{\mathcal{H}} + \|u\|_{\mathcal{H}},$$

where  $\|\nabla u\|_{\mathcal{H}} = \|\nabla u\|_{\mathcal{H}}$ . The completion of  $C_0^{\infty}(\Omega)$  in  $W^{1,\mathcal{H}}(\Omega)$  is denoted by  $W_0^{1,\mathcal{H}}(\Omega)$  and from (H)(i) we have an equivalent norm on  $W_0^{1,\mathcal{H}}(\Omega)$  given by

$$\|u\|_{1,\mathcal{H},0} = \|\nabla u\|_{\mathcal{H}},$$

see Proposition 2.18(ii) of Crespo-Blanco-Gasiński-Harjulehto-Winkert [17]. We know that  $W^{1,\mathcal{H}}(\Omega)$  and  $W_0^{1,\mathcal{H}}(\Omega)$  are reflexive Banach spaces.

We have the following embedding results for the spaces  $L^{\mathcal{H}}(\Omega)$  and  $W_0^{1,\mathcal{H}}(\Omega)$ .

**Proposition 2.1.** *Let (H)(i) be satisfied. Then the following embeddings hold:*

- (i)  $L^{\mathcal{H}}(\Omega) \hookrightarrow L^r(\Omega)$  and  $W_0^{1,\mathcal{H}}(\Omega) \hookrightarrow W_0^{1,r}(\Omega)$  are continuous for all  $r \in [1, p]$ ;
- (ii)  $W_0^{1,\mathcal{H}}(\Omega) \hookrightarrow L^r(\Omega)$  is continuous for all  $r \in [1, p^*]$ ;
- (iii)  $W_0^{1,\mathcal{H}}(\Omega) \hookrightarrow L^r(\Omega)$  is compact for all  $r \in [1, p^*]$ ;
- (iv)  $L^{\mathcal{H}}(\Omega) \hookrightarrow L_{\mu}^q(\Omega)$  is continuous;
- (v)  $L^q(\Omega) \hookrightarrow L^{\mathcal{H}}(\Omega)$  is continuous.

The norm  $\|\cdot\|_{\mathcal{H}}$  and the modular function  $\rho_{\mathcal{H}}$  are related as follows, see Liu-Dai [34, Proposition 2.1].

**Proposition 2.2.** *Let (H)(i) be satisfied, let  $y \in L^{\mathcal{H}}(\Omega)$  and let  $\rho_{\mathcal{H}}$  be defined by (2.1). Then the following hold:*

- (i) If  $y \neq 0$ , then  $\|y\|_{\mathcal{H}} = \lambda$  if and only if  $\rho_{\mathcal{H}}(\frac{y}{\lambda}) = 1$ ;
- (ii)  $\|y\|_{\mathcal{H}} < 1$  (resp.  $> 1, = 1$ ) if and only if  $\rho_{\mathcal{H}}(y) < 1$  (resp.  $> 1, = 1$ );
- (iii) If  $\|y\|_{\mathcal{H}} < 1$ , then  $\|y\|_{\mathcal{H}}^q \leq \rho_{\mathcal{H}}(y) \leq \|y\|_{\mathcal{H}}^p$ ;
- (iv) If  $\|y\|_{\mathcal{H}} > 1$ , then  $\|y\|_{\mathcal{H}}^p \leq \rho_{\mathcal{H}}(y) \leq \|y\|_{\mathcal{H}}^q$ ;
- (v)  $\|y\|_{\mathcal{H}} \rightarrow 0$  if and only if  $\rho_{\mathcal{H}}(y) \rightarrow 0$ ;
- (vi)  $\|y\|_{\mathcal{H}} \rightarrow +\infty$  if and only if  $\rho_{\mathcal{H}}(y) \rightarrow +\infty$ .

Let  $A: W_0^{1,\mathcal{H}}(\Omega) \rightarrow W_0^{1,\mathcal{H}}(\Omega)^*$  be the nonlinear map defined by

$$\langle A(u), \varphi \rangle_{\mathcal{H}} := \int_{\Omega} (|\nabla u|^{p-2} \nabla u + \mu(x) |\nabla u|^{q-2} \nabla u) \cdot \nabla \varphi dx \quad (2.2)$$

for all  $u, \varphi \in W_0^{1,\mathcal{H}}(\Omega)$ , where  $\langle \cdot, \cdot \rangle_{\mathcal{H}}$  is the duality pairing between  $W_0^{1,\mathcal{H}}(\Omega)$  and its dual space  $W_0^{1,\mathcal{H}}(\Omega)^*$ . The operator  $A: W_0^{1,\mathcal{H}}(\Omega) \rightarrow W_0^{1,\mathcal{H}}(\Omega)^*$  has the following properties, see Liu-Dai [34].

**Proposition 2.3.** *The operator  $A$  defined by (2.2) is bounded (that is, it maps bounded sets into bounded sets), continuous, strictly monotone (hence maximal monotone) and it is of type  $(S_+)$ .*

## 3. PROOF OF THE MAIN RESULT

In this section we are going to prove our main result stated as Theorem 1.1 in Section 1.

To this end, recall that  $\varphi_\lambda : W_0^{1,\mathcal{H}}(\Omega) \rightarrow \mathbb{R}$  is the corresponding energy function for problem (1.3) given by

$$\varphi_\lambda(u) = \frac{1}{p} \|\nabla u\|_p^p + \frac{1}{q} \|\nabla u\|_{q,\mu}^q - \frac{1}{1-\gamma} \int_\Omega a(x)|u|^{1-\gamma} dx - \frac{\lambda}{r} \|u\|_r^r.$$

Due to the presence of the singular term  $a(x)|u|^{1-\gamma}$  we know that  $\varphi_\lambda$  is not  $C^1$ . In order to overcome this, we will make use of the fibering method along with the Nehari manifold mentioned in the Introduction. Now we consider the fibering function  $\omega_u : [0, +\infty) \rightarrow \mathbb{R}$  for  $u \in W_0^{1,\mathcal{H}}(\Omega)$  defined by

$$\omega_u(t) = \varphi_\lambda(tu) \quad \text{for all } t \geq 0.$$

The Nehari manifold corresponding to the functional  $\varphi_\lambda$  is defined by

$$\begin{aligned} \mathcal{N}_\lambda &= \left\{ u \in W_0^{1,\mathcal{H}}(\Omega) \setminus \{0\} : \|\nabla u\|_p^p + \|\nabla u\|_{q,\mu}^q = \int_\Omega a(x)|u|^{1-\gamma} dx + \lambda \|u\|_r^r \right\} \\ &= \left\{ u \in W_0^{1,\mathcal{H}}(\Omega) \setminus \{0\} : \omega'_u(1) = 0 \right\}. \end{aligned}$$

It is easy to see that  $\mathcal{N}_\lambda$  is smaller than  $W_0^{1,\mathcal{H}}(\Omega)$  and it contains the weak solutions of problem (1.3). We will see that the functional  $\varphi_\lambda$  has nice properties restricted to  $\mathcal{N}_\lambda$  which fail globally. For our further considerations we need to decompose the set  $\mathcal{N}_\lambda$  in the following way:

$$\begin{aligned} \mathcal{N}_\lambda^+ &= \{ u \in \mathcal{N}_\lambda : (p + \gamma - 1)\|\nabla u\|_p^p + (q + \gamma - 1)\|\nabla u\|_{q,\mu}^q - \lambda(r + \gamma - 1)\|u\|_r^r > 0 \} \\ &= \{ u \in \mathcal{N}_\lambda : \omega''_u(1) > 0 \}, \\ \mathcal{N}_\lambda^0 &= \{ u \in \mathcal{N}_\lambda : (p + \gamma - 1)\|\nabla u\|_p^p + (q + \gamma - 1)\|\nabla u\|_{q,\mu}^q = \lambda(r + \gamma - 1)\|u\|_r^r \} \\ &= \{ u \in \mathcal{N}_\lambda : \omega''_u(1) = 0 \}, \\ \mathcal{N}_\lambda^- &= \{ u \in \mathcal{N}_\lambda : (p + \gamma - 1)\|\nabla u\|_p^p + (q + \gamma - 1)\|\nabla u\|_{q,\mu}^q - \lambda(r + \gamma - 1)\|u\|_r^r < 0 \} \\ &= \{ u \in \mathcal{N}_\lambda : \omega''_u(1) < 0 \}. \end{aligned}$$

We start with the following proposition about the coercivity of the energy functional  $\varphi_\lambda$  restricted to  $\mathcal{N}_\lambda$ .

**Proposition 3.1.** *Let hypotheses (H) be satisfied. Then  $\varphi_\lambda|_{\mathcal{N}_\lambda}$  is coercive.*

*Proof.* Let  $u \in \mathcal{N}_\lambda$  with  $\|u\|_{1,\mathcal{H},0} > 1$ . From the definition of the Nehari manifold  $\mathcal{N}_\lambda$  we have

$$-\frac{\lambda}{r} \|u\|_r^r = -\frac{1}{r} \|\nabla u\|_p^p - \frac{1}{r} \|\nabla u\|_{q,\mu}^q + \frac{1}{r} \int_\Omega a(x)|u|^{1-\gamma} dx. \quad (3.1)$$

Combining (3.1) with  $\varphi_\lambda$  and applying Proposition 2.2(iv) along with Theorem 13.17 of Hewitt-Stromberg [32, p. 196] gives

$$\begin{aligned} \varphi_\lambda(u) &= \left[ \frac{1}{p} - \frac{1}{r} \right] \|\nabla u\|_p^p + \left[ \frac{1}{q} - \frac{1}{r} \right] \|\nabla u\|_{q,\mu}^q + \left[ \frac{1}{r} - \frac{1}{1-\gamma} \right] \int_\Omega a(x)|u|^{1-\gamma} dx \\ &\geq \left[ \frac{1}{q} - \frac{1}{r} \right] \rho_{\mathcal{H}}(\nabla u) + \left[ \frac{1}{r} - \frac{1}{1-\gamma} \right] \int_\Omega a(x)|u|^{1-\gamma} dx \\ &\geq c_1 \|u\|_{1,\mathcal{H},0}^p - c_2 \|u\|_{1,\mathcal{H},0}^{1-\gamma} \end{aligned}$$

for some  $c_1, c_2 > 0$  because of  $p < q < r$ . Hence, due to  $1 - \gamma < 1 < p$ , the assertion of the proposition follows.  $\square$

Let  $m_\lambda^+ = \inf_{\mathcal{N}_\lambda^+} \varphi_\lambda$ .

**Proposition 3.2.** *Let hypotheses (H) be satisfied and suppose that  $\mathcal{N}_\lambda^+ \neq \emptyset$ . Then  $m_\lambda^+ < 0$ .*

*Proof.* Let  $u \in \mathcal{N}_\lambda^+$ . First note that  $\mathcal{N}_\lambda^+ \subseteq \mathcal{N}_\lambda$  which implies that

$$-\frac{1}{1-\gamma} \int_{\Omega} a(x)|u|^{1-\gamma} dx = -\frac{1}{1-\gamma} (\|\nabla u\|_p^p + \|\nabla u\|_{q,\mu}^q) + \frac{\lambda}{1-\gamma} \|u\|_r^r. \quad (3.2)$$

On the other hand, by definition of  $\mathcal{N}_\lambda^+$ , we have

$$\lambda \|u\|_r^r < \frac{p+\gamma-1}{r+\gamma-1} \|\nabla u\|_p^p + \frac{q+\gamma-1}{r+\gamma-1} \|\nabla u\|_{q,\mu}^q. \quad (3.3)$$

From (3.3) and (3.2) it follows that

$$\begin{aligned} \varphi_\lambda(u) &= \frac{1}{p} \|\nabla u\|_p^p + \frac{1}{q} \|\nabla u\|_{q,\mu}^q - \frac{1}{1-\gamma} \int_{\Omega} a(x)|u|^{1-\gamma} dx - \frac{\lambda}{r} \|u\|_r^r \\ &= \left[ \frac{1}{p} - \frac{1}{1-\gamma} \right] \|\nabla u\|_p^p + \left[ \frac{1}{q} - \frac{1}{1-\gamma} \right] \|\nabla u\|_{q,\mu}^q + \lambda \left[ \frac{1}{1-\gamma} - \frac{1}{r} \right] \|u\|_r^r \\ &\leq \left[ \frac{-(p+\gamma-1)}{p(1-\gamma)} + \frac{p+\gamma-1}{r+\gamma-1} \cdot \frac{r+\gamma-1}{r(1-\gamma)} \right] \|\nabla u\|_p^p \\ &\quad + \left[ \frac{-(q+\gamma-1)}{q(1-\gamma)} + \frac{q+\gamma-1}{r+\gamma-1} \cdot \frac{r+\gamma-1}{r(1-\gamma)} \right] \|\nabla u\|_{q,\mu}^q \\ &= \frac{p+\gamma-1}{1-\gamma} \left[ \frac{1}{r} - \frac{1}{p} \right] \|\nabla u\|_p^p + \frac{q+\gamma-1}{1-\gamma} \left[ \frac{1}{r} - \frac{1}{q} \right] \|\nabla u\|_{q,\mu}^q \\ &< 0, \end{aligned}$$

since  $p < q < r$ . Hence,  $\varphi_\lambda|_{\mathcal{N}_\lambda^+} < 0$  and so  $m_\lambda^+ < 0$ .  $\square$

**Proposition 3.3.** *Let hypotheses (H) be satisfied. Then there exists  $\lambda^* > 0$  such that  $\mathcal{N}_\lambda^0 = \emptyset$  for all  $\lambda \in (0, \lambda^*)$ .*

*Proof.* Arguing indirectly, suppose that for every  $\lambda^* > 0$  there exists  $\lambda \in (0, \lambda^*)$  such that  $\mathcal{N}_\lambda^0 \neq \emptyset$ . Hence, for any given  $\lambda > 0$ , we can find  $u \in \mathcal{N}_\lambda^0$  such that

$$(p+\gamma-1)\|\nabla u\|_p^p + (q+\gamma-1)\|\nabla u\|_{q,\mu}^q = \lambda(r+\gamma-1)\|u\|_r^r. \quad (3.4)$$

Since  $u \in \mathcal{N}_\lambda$ , one also has

$$\begin{aligned} &(r+\gamma-1)\|\nabla u\|_p^p + (r+\gamma-1)\|\nabla u\|_{q,\mu}^q \\ &= (r+\gamma-1) \int_{\Omega} a(x)|u|^{1-\gamma} dx + \lambda(r+\gamma-1)\|u\|_r^r. \end{aligned} \quad (3.5)$$

Subtracting (3.4) from (3.5) yields

$$(r-p)\|\nabla u\|_p^p + (r-q)\|\nabla u\|_{q,\mu}^q = (r+\gamma-1) \int_{\Omega} a(x)|u|^{1-\gamma} dx. \quad (3.6)$$

Applying Proposition 2.2(iii), (iv), Theorem 13.17 of Hewitt-Stromberg [32, p. 196] and Proposition 2.1(ii) we get from (3.6) that

$$\min \left\{ \|u\|_{1,\mathcal{H},0}^p, \|u\|_{1,\mathcal{H},0}^q \right\} \leq c_3 \|u\|_{1,\mathcal{H},0}^{1-\gamma}$$

for some  $c_3 > 0$  since  $1 - \gamma < 1 < p < q < r$ . Hence

$$\|u\|_{1,\mathcal{H},0} \leq c_4 \quad (3.7)$$

for some  $c_4 > 0$ .

On the other hand, from (3.4), Proposition 2.2(iii), (iv) and Proposition 2.1(ii) we have

$$\min \left\{ \|u\|_{1,\mathcal{H},0}^p, \|u\|_{1,\mathcal{H},0}^q \right\} \leq \lambda c_5 \|u\|_{1,\mathcal{H},0}^r$$

for some  $c_5 > 0$ . Consequently,

$$\|u\|_{1,\mathcal{H},0} \geq \left( \frac{1}{\lambda c_5} \right)^{\frac{1}{r-p}} \quad \text{or} \quad \|u\|_{1,\mathcal{H},0} \geq \left( \frac{1}{\lambda c_5} \right)^{\frac{1}{r-q}}.$$

If  $\lambda \rightarrow 0^+$ , due to  $p < q < r$ , then  $\|u\|_{1,\mathcal{H},0} \rightarrow +\infty$ , which contradicts (3.7).  $\square$

**Proposition 3.4.** *Let hypotheses (H) be satisfied. Then there exists  $\hat{\lambda}^* \in (0, \lambda^*]$  such that  $\mathcal{N}_\lambda^\pm \neq \emptyset$  for all  $\lambda \in (0, \hat{\lambda}^*)$ . In addition, for any  $\lambda \in (0, \hat{\lambda}^*)$ , there exists  $u^* \in \mathcal{N}_\lambda^+$  such that  $\varphi_\lambda(u^*) = m_\lambda^+ < 0$  and  $u^*(x) \geq 0$  for a. a.  $x \in \Omega$ .*

*Proof.* Let  $u \in W_0^{1,\mathcal{H}}(\Omega) \setminus \{0\}$  and consider the function  $\hat{\psi}_u: (0, +\infty) \rightarrow \mathbb{R}$  defined by

$$\hat{\psi}_u(t) = t^{p-r} \|\nabla u\|_p^p - t^{-r-\gamma+1} \int_\Omega a(x) |u|^{1-\gamma} dx.$$

Since  $r - p < r + \gamma - 1$  we can find  $\hat{t}_0 > 0$  such that

$$\hat{\psi}_u(\hat{t}_0) = \max_{t>0} \hat{\psi}_u(t).$$

Thus,  $\hat{\psi}'_u(\hat{t}_0) = 0$ , that is,

$$(p-r)\hat{t}_0^{p-r-1} \|\nabla u\|_p^p + (r+\gamma-1)\hat{t}_0^{-r-\gamma} \int_\Omega a(x) |u|^{1-\gamma} dx = 0.$$

Hence

$$\hat{t}_0 = \left[ \frac{(r+\gamma-1) \int_\Omega a(x) |u|^{1-\gamma} dx}{(r-p) \|\nabla u\|_p^p} \right]^{\frac{1}{p+\gamma-1}}.$$

Moreover, we have

$$\begin{aligned} \hat{\psi}_u(\hat{t}_0) &= \frac{\left[ (r-p) \|\nabla u\|_p^p \right]^{\frac{r-p}{p+\gamma-1}}}{\left[ (r+\gamma-1) \int_\Omega a(x) |u|^{1-\gamma} dx \right]^{\frac{r-p}{p+\gamma-1}}} \|\nabla u\|_p^p \\ &\quad - \frac{\left[ (r-p) \|\nabla u\|_p^p \right]^{\frac{r+\gamma-1}{p+\gamma-1}}}{\left[ (r+\gamma-1) \int_\Omega a(x) |u|^{1-\gamma} dx \right]^{\frac{r+\gamma-1}{p+\gamma-1}}} \int_\Omega a(x) |u|^{1-\gamma} dx \\ &= \frac{(r-p) \|\nabla u\|_p^{\frac{p(r+\gamma-1)}{p+\gamma-1}}}{(r+\gamma-1) \int_\Omega a(x) |u|^{1-\gamma} dx} \end{aligned} \quad (3.8)$$

$$\begin{aligned}
& - \frac{(r-p)^{\frac{r+\gamma-1}{p+\gamma-1}} \|\nabla u\|_p^{\frac{p(r+\gamma-1)}{p+\gamma-1}}}{(r+\gamma-1)^{\frac{r+\gamma-1}{p+\gamma-1}} \left[ \int_{\Omega} a(x)|u|^{1-\gamma} dx \right]^{\frac{r-p}{p+\gamma-1}}} \\
& = \frac{p+\gamma-1}{r-p} \left[ \frac{r-p}{r+\gamma-1} \right]^{\frac{r+\gamma-1}{p+\gamma-1}} \frac{\|\nabla u\|_p^{\frac{p(r+\gamma-1)}{p+\gamma-1}}}{\left[ \int_{\Omega} a(x)|u|^{1-\gamma} dx \right]^{\frac{r-p}{p+\gamma-1}}}.
\end{aligned}$$

Let  $S$  be the best constant of the Sobolev embedding  $W_0^{1,p}(\Omega) \rightarrow L^{p^*}(\Omega)$ , that is,

$$S\|u\|_{p^*}^p \leq \|\nabla u\|_p^p. \quad (3.9)$$

Moreover, we have

$$\int_{\Omega} a(x)|u|^{1-\gamma} dx \leq c_6 \|u\|_{p^*}^{1-\gamma} \quad (3.10)$$

for some  $c_6 > 0$ . Combining (3.8), (3.9) and (3.10) gives

$$\begin{aligned}
& \hat{\psi}_u(\hat{t}_0) - \lambda \|u\|_r^r \\
& = \frac{p+\gamma-1}{r-p} \left[ \frac{r-p}{r+\gamma-1} \right]^{\frac{r+\gamma-1}{p+\gamma-1}} \frac{\|\nabla u\|_p^{\frac{p(r+\gamma-1)}{p+\gamma-1}}}{\left[ \int_{\Omega} a(x)|u|^{1-\gamma} dx \right]^{\frac{r-p}{p+\gamma-1}}} - \lambda \|u\|_r^r \\
& \geq \frac{p+\gamma-1}{r-p} \left[ \frac{r-p}{r+\gamma-1} \right]^{\frac{r+\gamma-1}{p+\gamma-1}} \frac{S^{\frac{r+\gamma-1}{p+\gamma-1}} (\|u\|_{p^*}^p)^{\frac{r+\gamma-1}{p+\gamma-1}}}{\left( c_6 \|u\|_{p^*}^{1-\gamma} \right)^{\frac{r-p}{p+\gamma-1}}} - \lambda c_7 \|u\|_{p^*}^r \\
& = [c_8 - \lambda c_7] \|u\|_{p^*}^r
\end{aligned}$$

for some  $c_7, c_8 > 0$ . Therefore, there exists  $\hat{\lambda}^* \in (0, \lambda^*]$  independent of  $u$  such that

$$\hat{\psi}_u(\hat{t}_0) - \lambda \|u\|_r^r > 0 \quad \text{for all } \lambda \in (0, \hat{\lambda}^*). \quad (3.11)$$

Now consider the function  $\psi_u: (0, +\infty) \rightarrow \mathbb{R}$  defined by

$$\psi_u(t) = t^{p-r} \|\nabla u\|_p^p + t^{q-r} \|\nabla u\|_{q,\mu}^q - t^{-r-\gamma+1} \int_{\Omega} a(x)|u|^{1-\gamma} dx.$$

Since  $r-q < r-p < r+\gamma-1$  we can find  $t_0 > 0$  such that

$$\psi_u(t_0) = \max_{t>0} \psi_u(t).$$

Because of  $\psi_u \geq \hat{\psi}_u$  and due to (3.11) (note that there the choice of  $\hat{\lambda}^*$  is independent of  $u$ ) we can find  $\hat{\lambda}^* \in (0, \lambda^*]$  independent of  $u$  such that

$$\psi_u(t_0) - \lambda \|u\|_r^r > 0 \quad \text{for all } \lambda \in (0, \hat{\lambda}^*).$$

Thus there exist  $t_1 < t_0 < t_2$  such that

$$\psi_u(t_1) = \lambda \|u\|_r^r = \psi_u(t_2) \quad \text{and} \quad \psi'_u(t_2) < 0 < \psi'_u(t_1), \quad (3.12)$$

where

$$\begin{aligned}
\psi'_u(t) & = (p-r)t^{p-r-1} \|\nabla u\|_p^p + (q-r)t^{q-r-1} \|\nabla u\|_{q,\mu}^q \\
& \quad - (-r-\gamma+1)t^{-r-\gamma} \int_{\Omega} a(x)|u|^{1-\gamma} dx.
\end{aligned} \quad (3.13)$$

Note that  $t_1, t_2$  are the only numbers which fulfill the equality in (3.12).

Recall that the fibering function  $\omega_u: [0, +\infty) \rightarrow \mathbb{R}$  is given by

$$\omega_u(t) = \varphi_\lambda(tu) \quad \text{for all } t \geq 0.$$

Clearly,  $\omega_u \in C^\infty((0, \infty))$ . We have

$$\omega'_u(t_1) = t_1^{p-1} \|\nabla u\|_p^p + t_1^{q-1} \|\nabla u\|_{q,\mu}^q - t_1^{-\gamma} \int_\Omega a(x)|u|^{1-\gamma} dx - \lambda t_1^{r-1} \|u\|_r^r$$

and

$$\begin{aligned} \omega''_u(t_1) &= (p-1)t_1^{p-2} \|\nabla u\|_p^p + (q-1)t_1^{q-2} \|\nabla u\|_{q,\mu}^q \\ &\quad + \gamma t_1^{-\gamma-1} \int_\Omega a(x)|u|^{1-\gamma} dx - \lambda(r-1)t_1^{r-2} \|u\|_r^r. \end{aligned} \quad (3.14)$$

From (3.12) we obtain

$$t_1^{p-r} \|\nabla u\|_p^p + t_1^{q-r} \|\nabla u\|_{q,\mu}^q - t_1^{-r-\gamma+1} \int_\Omega a(x)|u|^{1-\gamma} dx = \lambda \|u\|_r^r,$$

which implies by multiplying with  $\gamma t_1^{r-2}$  and  $-(r-1)t_1^{r-2}$ , respectively, that

$$\gamma t_1^{p-2} \|\nabla u\|_p^p + \gamma t_1^{q-2} \|\nabla u\|_{q,\mu}^q - \gamma \lambda t_1^{r-2} \|u\|_r^r = \gamma t_1^{-\gamma-1} \int_\Omega a(x)|u|^{1-\gamma} dx \quad (3.15)$$

and

$$\begin{aligned} &-(r-1)t_1^{p-2} \|\nabla u\|_p^p - (r-1)t_1^{q-2} \|\nabla u\|_{q,\mu}^q \\ &\quad + (r-1)t_1^{-\gamma-1} \int_\Omega a(x)|u|^{1-\gamma} dx \\ &= -\lambda(r-1)t_1^{r-2} \|u\|_r^r. \end{aligned} \quad (3.16)$$

Applying (3.15) in (3.14) gives

$$\begin{aligned} \omega''_u(t_1) &= (p+\gamma-1)t_1^{p-2} \|\nabla u\|_p^p + (q+\gamma-1)t_1^{q-2} \|\nabla u\|_{q,\mu}^q \\ &\quad - \lambda(r+\gamma-1)t_1^{r-2} \|u\|_r^r \\ &= t_1^{-2} \left[ (p+\gamma-1)t_1^p \|\nabla u\|_p^p + (q+\gamma-1)t_1^q \|\nabla u\|_{q,\mu}^q \right. \\ &\quad \left. - \lambda(r+\gamma-1)t_1^r \|u\|_r^r \right]. \end{aligned} \quad (3.17)$$

On the other hand, applying (3.16) in (3.14) and using the representation in (3.13) leads to

$$\begin{aligned} \omega''_u(t_1) &= (p-r)t_1^{p-2} \|\nabla u\|_p^p + (q-r)t_1^{q-2} \|\nabla u\|_{q,\mu}^q \\ &\quad + (r+\gamma-1)t_1^{-\gamma-1} \int_\Omega a(x)|u|^{1-\gamma} dx \\ &= t_1^{1-r} \psi'_u(t_1) > 0. \end{aligned} \quad (3.18)$$

From (3.17) and (3.18) it follows that

$$(p+\gamma-1)t_1^p \|\nabla u\|_p^p + (q+\gamma-1)t_1^q \|\nabla u\|_{q,\mu}^q - \lambda(r+\gamma-1)t_1^r \|u\|_r^r > 0,$$

which implies

$$t_1 u \in \mathcal{N}_\lambda^+ \quad \text{for all } \lambda \in (0, \hat{\lambda}^*].$$

Hence,  $\mathcal{N}_\lambda^+ \neq \emptyset$ .

Using similar arguments for the point  $t_2$  (see (3.12)), we can show that  $\mathcal{N}_\lambda^- \neq \emptyset$ . This shows the first assertion of the proposition. Let us now prove the second one.

To this end, let  $\{u_n\}_{n \in \mathbb{N}} \subset \mathcal{N}_\lambda^+$  be a minimizing sequence, that is,

$$\varphi_\lambda(u_n) \searrow m_\lambda^+ < 0 \quad \text{as } n \rightarrow \infty. \quad (3.19)$$

Recall that  $\mathcal{N}_\lambda^+ \subset \mathcal{N}_\lambda$  and so we conclude from Proposition 3.1 that  $\{u_n\}_{n \in \mathbb{N}} \subset W_0^{1,\mathcal{H}}(\Omega)$  is bounded. Therefore, we may assume that

$$u_n \rightharpoonup u^* \quad \text{in } W_0^{1,\mathcal{H}}(\Omega) \quad \text{and} \quad u_n \rightarrow u^* \quad \text{in } L^r(\Omega). \quad (3.20)$$

From (3.19) and (3.20) we know that

$$\varphi_\lambda(u^*) \leq \liminf_{n \rightarrow +\infty} \varphi_\lambda(u_n) < 0 = \varphi_\lambda(0).$$

Hence,  $u^* \neq 0$ .

**Claim:**  $\liminf_{n \rightarrow +\infty} \rho_{\mathcal{H}}(u_n) = \rho_{\mathcal{H}}(u^*)$

Suppose, by contradiction, that

$$\liminf_{n \rightarrow +\infty} \rho_{\mathcal{H}}(u_n) > \rho_{\mathcal{H}}(u^*).$$

Then, by using (3.12), we have

$$\begin{aligned} & \liminf_{n \rightarrow +\infty} \omega'_{u_n}(t_1) \\ &= \liminf_{n \rightarrow +\infty} \left[ t_1^{p-1} \|\nabla u_n\|_p^p + t_1^{q-1} \|\nabla u_n\|_{q,\mu}^q - t_1^{-\gamma} \int_{\Omega} a(x) |u_n|^{1-\gamma} dx - \lambda t_1^{r-1} \|u_n\|_r^r \right] \\ &> t_1^{p-1} \|\nabla u^*\|_p^p + t_1^{q-1} \|\nabla u^*\|_{q,\mu}^q - t_1^{-\gamma} \int_{\Omega} a(x) |u^*|^{1-\gamma} dx - \lambda t_1^{r-1} \|u^*\|_r^r \\ &= \omega'_{u^*}(t_1) = t_1^{r-1} [\psi_{u^*}(t_1) - \lambda \|u^*\|_r^r] = 0, \end{aligned}$$

which implies the existence of  $n_0 \in \mathbb{N}$  such that  $\omega'_{u_n}(t_1) > 0$  for all  $n > n_0$ . Recall that  $u_n \in \mathcal{N}_\lambda^+ \subset \mathcal{N}_\lambda$  and  $\omega'_{u_n}(t) = t^{r-1} [\psi_{u_n}(t) - \lambda \|u_n\|_r^r]$ . Thus we have  $\omega'_{u_n}(t) < 0$  for all  $t \in (0, 1)$  and  $\omega'_{u_n}(1) = 0$ . Therefore,  $t_1 > 1$ .

Since  $\omega_{u^*}$  is decreasing on  $(0, t_1]$ , we have

$$\varphi_\lambda(t_1 u^*) \leq \varphi_\lambda(u^*) < m_\lambda^+.$$

Recall that  $t_1 u^* \in \mathcal{N}_\lambda^+$ . So we obtain that

$$m_\lambda^+ \leq \varphi_\lambda(t_1 u^*) < m_\lambda^+,$$

a contradiction. So the Claim is proved.

From the Claim we know that we can find a subsequence (still denoted by  $u_n$ ) such that  $\rho_{\mathcal{H}}(u_n) \rightarrow \rho_{\mathcal{H}}(u^*)$ . It follows from Proposition 2.2(v) that  $u_n \rightarrow u$  in  $W_0^{1,\mathcal{H}}(\Omega)$ . This implies  $\varphi_\lambda(u_n) \rightarrow \varphi_\lambda(u^*)$ , and consequently,  $\varphi_\lambda(u^*) = m_\lambda^+$ . Since  $u_n \in \mathcal{N}_\lambda^+$  for all  $n \in \mathbb{N}$ , we have

$$(p + \gamma - 1) \|\nabla u_n\|_p^p + (q + \gamma - 1) \|\nabla u_n\|_{q,\mu}^q - \lambda(r + \gamma - 1) \|u_n\|_r^r > 0.$$

Letting  $n \rightarrow +\infty$  gives

$$(p + \gamma - 1) \|\nabla u^*\|_p^p + (q + \gamma - 1) \|\nabla u^*\|_{q,\mu}^q - \lambda(r + \gamma - 1) \|u^*\|_r^r \geq 0. \quad (3.21)$$

Recall that  $\lambda \in (0, \hat{\lambda}^*)$  and  $\hat{\lambda}^* \leq \lambda^*$ . Then, from Proposition 3.3 we know that equality in (3.21) cannot occur. Therefore, we conclude that  $u^* \in \mathcal{N}_\lambda^+$ . Since we can use  $|u^*|$  instead of  $u^*$ , we may assume that  $u^*(x) \geq 0$  for a. a.  $x \in \Omega$  with  $u^* \neq 0$ . The proof is finished.  $\square$

In what follows, for  $\varepsilon > 0$ , we denote

$$B_\varepsilon(0) = \left\{ u \in W_0^{1,\mathcal{H}}(\Omega) : \|u\|_{1,\mathcal{H},0} < \varepsilon \right\}.$$

The next lemma is motivated by Lemma 3 of Sun-Wu-Long [56]. This lemma is helpful in order to show that  $u^*$  is a local minimizer of  $\varphi_\lambda$  (see Proposition 3.6) and from this we conclude that  $u^*$  is a weak solution of (1.3) (see Proposition 3.7).

**Lemma 3.5.** *Let hypotheses (H) be satisfied and let  $u \in \mathcal{N}_\lambda^\pm$ . Then there exist  $\varepsilon > 0$  and a continuous function  $\vartheta: B_\varepsilon(0) \rightarrow (0, \infty)$  such that*

$$\vartheta(0) = 1 \quad \text{and} \quad \vartheta(y)(u + y) \in \mathcal{N}_\lambda^\pm \quad \text{for all } y \in B_\varepsilon(0).$$

*Proof.* We show the proof only for  $\mathcal{N}_\lambda^+$ , the proof for  $\mathcal{N}_\lambda^-$  works in a similar way. To this end, let  $\zeta: W_0^{1,\mathcal{H}}(\Omega) \times (0, \infty) \rightarrow \mathbb{R}$  be defined by

$$\begin{aligned} \zeta(y, t) &= t^{p+\gamma-1} \|\nabla(u+y)\|_p^p + t^{q+\gamma-1} \|\nabla(u+y)\|_{q,\mu}^q - \int_\Omega a(x) |u+y|^{1-\gamma} dx \\ &\quad - \lambda t^{r+\gamma-1} \|u+y\|_r^r \quad \text{for all } y \in W_0^{1,\mathcal{H}}(\Omega). \end{aligned}$$

Since  $u \in \mathcal{N}_\lambda^+ \subset \mathcal{N}_\lambda$ , one has  $\zeta(0, 1) = 0$ . Because of  $u \in \mathcal{N}_\lambda^+$ , it follows that

$$\zeta'_t(0, 1) = (p + \gamma - 1) \|\nabla u\|_p^p + (q + \gamma - 1) \|\nabla u\|_{q,\mu}^q - \lambda(r + \gamma - 1) \|u\|_r^r > 0.$$

Then, by the implicit function theorem, see, for example, Gasiński-Papageorgiou [27, p. 481], there exist  $\varepsilon > 0$  and a continuous function  $\vartheta: B_\varepsilon(0) \rightarrow (0, \infty)$  such that

$$\vartheta(0) = 1 \quad \text{and} \quad \vartheta(y)(u + y) \in \mathcal{N}_\lambda \quad \text{for all } y \in B_\varepsilon(0).$$

Choosing  $\varepsilon > 0$  small enough, we also have

$$\vartheta(0) = 1 \quad \text{and} \quad \vartheta(y)(u + y) \in \mathcal{N}_\lambda^+ \quad \text{for all } y \in B_\varepsilon(0).$$

□

**Proposition 3.6.** *Let hypotheses (H) be satisfied, let  $h \in W_0^{1,\mathcal{H}}(\Omega)$  and let  $\lambda \in (0, \hat{\lambda}^*]$ . Then there exists  $b > 0$  such that  $\varphi_\lambda(u^*) \leq \varphi_\lambda(u^* + th)$  for all  $t \in [0, b]$ .*

*Proof.* We introduce the function  $\eta_h: [0, +\infty) \rightarrow \mathbb{R}$  defined by

$$\begin{aligned} \eta_h(t) &= (p-1) \|\nabla u^* + t\nabla h\|_p^p + (q-1) \|\nabla u^* + t\nabla h\|_{q,\mu}^q \\ &\quad + \gamma \int_\Omega a(x) |u^* + th|^{1-\gamma} dx - \lambda(r-1) \|u^* + th\|_r^r. \end{aligned} \quad (3.22)$$

Recall that  $u^* \in \mathcal{N}_\lambda^+ \subseteq \mathcal{N}_\lambda$ , see Proposition 3.4. This implies

$$\gamma \int_\Omega a(x) |u^*|^{1-\gamma} dx = \gamma \|\nabla u^*\|_p^p + \gamma \|\nabla u^*\|_{q,\mu}^q - \lambda\gamma \|u^*\|_r^r \quad (3.23)$$

and

$$(p + \gamma - 1) \|\nabla u^*\|_p^p + (q + \gamma - 1) \|\nabla u^*\|_{q,\mu}^q - \lambda(r + \gamma - 1) \|u^*\|_r^r > 0. \quad (3.24)$$

Combining (3.22), (3.23) and (3.24) we see that  $\eta_h(0) > 0$ . Since  $\eta_h: [0, +\infty) \rightarrow \mathbb{R}$  is continuous we can find  $b_0 > 0$  such that

$$\eta_h(t) > 0 \quad \text{for all } t \in [0, b_0].$$

Lemma 3.5 implies that for every  $t \in [0, b_0]$  we can find  $\vartheta(t) > 0$  such that

$$\vartheta(t)(u^* + th) \in \mathcal{N}_\lambda^+ \quad \text{and} \quad \vartheta(t) \rightarrow 1 \quad \text{as } t \rightarrow 0^+. \quad (3.25)$$

From Proposition 3.4 we know that

$$m_\lambda^+ = \varphi_\lambda(u^*) \leq \varphi_\lambda(\vartheta(t)(u^* + th)) \quad \text{for all } t \in [0, b_0]. \quad (3.26)$$

From  $\omega_{u^*}''(1) > 0$  and the continuity in  $t$ , we have  $\omega_{u^*+th}''(1) > 0$  for  $t \in [0, b]$  with  $b \in (0, b_0]$ . Combining this with (3.26) gives

$$m_\lambda^+ = \varphi_\lambda(u^*) \leq \varphi_\lambda(\vartheta(t)(u^* + th)) = \omega_{u^*+th}(\vartheta(t)) \leq \omega_{u^*+th}(1) = \varphi_\lambda(u^* + th)$$

for all  $t \in [0, b]$ .  $\square$

The next proposition shows that  $\mathcal{N}_\lambda^+$  is a natural constraint for the energy functional  $\varphi_\lambda$ , see Papageorgiou-Rădulescu-Repovš [48, p. 425].

**Proposition 3.7.** *Let hypotheses (H) be satisfied and let  $\lambda \in (0, \hat{\lambda}^*]$ . Then  $u^*$  is a weak solution of problem (1.3) such that  $\varphi_\lambda(u^*) < 0$ .*

*Proof.* From Proposition 3.4 we know that  $u^* \geq 0$  for a. a.  $x \in \Omega$  and  $\varphi_\lambda(u^*) < 0$ .

Let us prove that  $u^* > 0$  for a. a.  $x \in \Omega$ . We argue indirectly and suppose there is a set  $D$  with positive measure such that  $u^*(x) = 0$  for a. a.  $x \in D$ . Now let  $h \in W_0^{1,\mathcal{H}}(\Omega)$  with  $h > 0$  and let  $t \in (0, b)$ , where  $b$  is from Proposition 3.6. Then we have  $(u^* + th)^{1-\gamma} > (u^*)^{1-\gamma}$  for a. a.  $x \in D$ . Applying this fact along with Proposition 3.6 results in

$$\begin{aligned} 0 &\leq \frac{\varphi_\lambda(u^* + th) - \varphi_\lambda(u^*)}{t} \\ &= \frac{1}{p} \frac{\|\nabla(u^* + th)\|_p^p - \|\nabla u^*\|_p^p}{t} + \frac{1}{q} \frac{\|\nabla(u^* + th)\|_{q,\mu}^q - \|\nabla u^*\|_{q,\mu}^q}{t} \\ &\quad - \frac{1}{(1-\gamma)t^\gamma} \int_D a(x)h^{1-\gamma} dx - \frac{1}{1-\gamma} \int_{\Omega \setminus D} a(x) \frac{(u^* + th)^{1-\gamma} - (u^*)^{1-\gamma}}{t} dx \\ &\quad - \frac{\lambda}{r} \frac{\|u^* + th\|_r^r - \|u^*\|_r^r}{t} \\ &< \frac{1}{p} \frac{\|\nabla(u^* + th)\|_p^p - \|\nabla u^*\|_p^p}{t} + \frac{1}{q} \frac{\|\nabla(u^* + th)\|_{q,\mu}^q - \|\nabla u^*\|_{q,\mu}^q}{t} \\ &\quad - \frac{1}{(1-\gamma)t^\gamma} \int_D a(x)h^{1-\gamma} dx - \frac{\lambda}{r} \frac{\|u^* + th\|_r^r - \|u^*\|_r^r}{t}. \end{aligned}$$

Since  $a > 0$ , see hypothesis (H)(iii), we conclude from the estimate above that

$$0 \leq \frac{\varphi_\lambda(u^* + th) - \varphi_\lambda(u^*)}{t} \rightarrow -\infty \quad \text{as } t \rightarrow 0^+.$$

This is a contradiction and so we have that  $u^*(x) > 0$  for a. a.  $x \in \Omega$ .

Next we prove that

$$a(\cdot)(u^*)^{-\gamma}h \in L^1(\Omega) \quad \text{for all } h \in W_0^{1,\mathcal{H}}(\Omega) \quad (3.27)$$

and

$$\begin{aligned} &\int_\Omega \left( |\nabla u^*|^{p-2} \nabla u^* + \mu(x) |\nabla u^*|^{q-2} \nabla u^* \right) \cdot \nabla h dx \\ &\geq \int_\Omega a(x)(u^*)^{-\gamma}h dx + \lambda \int_\Omega (u^*)^{r-1}h dx \end{aligned} \quad (3.28)$$

for all  $h \in W_0^{1,\mathcal{H}}(\Omega)$  with  $h \geq 0$ .

To this end, let  $h \in W_0^{1,\mathcal{H}}(\Omega)$  with  $h \geq 0$  and let  $\{t_n\}_{n \in \mathbb{N}} \subseteq (0, 1]$  be a decreasing sequence such that  $\lim_{n \rightarrow \infty} t_n = 0$ . First note that the functions

$$\kappa_n(x) = a(x) \frac{(u^*(x) + t_n h(x))^{1-\gamma} - u^*(x)^{1-\gamma}}{t_n}, \quad n \in \mathbb{N}$$

are nonnegative and measurable. Furthermore, we have

$$\lim_{n \rightarrow \infty} \kappa_n(x) = (1 - \gamma) a(x) u^*(x)^{-\gamma} h(x) \quad \text{for a. a. } x \in \Omega$$

and by Fatou's lemma we get

$$\int_{\Omega} a(x) (u^*)^{-\gamma} h \, dx \leq \frac{1}{1 - \gamma} \liminf_{n \rightarrow \infty} \int_{\Omega} \kappa_n \, dx. \quad (3.29)$$

Again from Proposition 3.6 we get for  $n \in \mathbb{N}$  sufficiently large that

$$\begin{aligned} 0 &\leq \frac{\varphi_{\lambda}(u^* + t_n h) - \varphi_{\lambda}(u^*)}{t_n} \\ &= \frac{1}{p} \frac{\|\nabla(u^* + t_n h)\|_p^p - \|\nabla u^*\|_p^p}{t_n} + \frac{1}{q} \frac{\|\nabla(u^* + t_n h)\|_{q,\mu}^q - \|\nabla u^*\|_{q,\mu}^q}{t_n} \\ &\quad - \frac{1}{1 - \gamma} \int_{\Omega} \kappa_n \, dx - \frac{\lambda}{r} \frac{\|u^* + t_n h\|_r^r - \|u^*\|_r^r}{t_n}. \end{aligned}$$

If we pass to the limit as  $n \rightarrow \infty$ , taking (3.29) into account, we obtain

$$\begin{aligned} &\int_{\Omega} a(x) (u^*)^{-\gamma} h \, dx \\ &\leq \int_{\Omega} (|\nabla u^*|^{p-2} \nabla u^* + \mu(x) |\nabla u^*|^{q-2} \nabla u^*) \cdot \nabla h \, dx - \lambda \int_{\Omega} (u^*)^{r-1} h \, dx. \end{aligned}$$

This shows (3.27) and (3.28). We point out that it is sufficient to prove the integrability in (3.27) for nonnegative test functions  $h \in W_0^{1,\mathcal{H}}(\Omega)$ .

In the next step we prove that  $u^*$  is a weak solution of (1.3). Let  $v \in W_0^{1,\mathcal{H}}(\Omega)$  and let  $\varepsilon > 0$ . We take  $h = (u^* + \varepsilon v)_+$  as test function in (3.28) and use  $u^* \in \mathcal{N}_{\lambda}^+ \subset \mathcal{N}_{\lambda}$  with  $u^* \geq 0$ . One has

$$\begin{aligned} 0 &\leq \int_{\Omega} (|\nabla u^*|^{p-2} \nabla u^* + \mu(x) |\nabla u^*|^{q-2} \nabla u^*) \cdot \nabla (u^* + \varepsilon v)_+ \, dx \\ &\quad - \int_{\Omega} (a(x) (u^*)^{-\gamma} + \lambda (u^*)^{r-1}) (u^* + \varepsilon v)_+ \, dx \\ &= \int_{\{u^* + \varepsilon v \geq 0\}} (|\nabla u^*|^{p-2} \nabla u^* + \mu(x) |\nabla u^*|^{q-2} \nabla u^*) \cdot \nabla (u^* + \varepsilon v) \, dx \\ &\quad - \int_{\{u^* + \varepsilon v \geq 0\}} (a(x) (u^*)^{-\gamma} + \lambda (u^*)^{r-1}) (u^* + \varepsilon v) \, dx \\ &= \int_{\Omega} (|\nabla u^*|^{p-2} \nabla u^* + \mu(x) |\nabla u^*|^{q-2} \nabla u^*) \cdot \nabla (u^* + \varepsilon v) \, dx \\ &\quad - \int_{\{u^* + \varepsilon v < 0\}} (|\nabla u^*|^{p-2} \nabla u^* + \mu(x) |\nabla u^*|^{q-2} \nabla u^*) \cdot \nabla (u^* + \varepsilon v) \, dx \\ &\quad - \int_{\Omega} (a(x) (u^*)^{-\gamma} + \lambda (u^*)^{r-1}) (u^* + \varepsilon v) \, dx \end{aligned}$$

$$\begin{aligned}
& + \int_{\{u^* + \varepsilon v < 0\}} \left( a(x) (u^*)^{-\gamma} + \lambda (u^*)^{r-1} \right) (u^* + \varepsilon v) \, dx \\
= & \|\nabla u^*\|_p^p + \|\nabla u^*\|_{q,\mu}^q - \int_{\Omega} a(x) |u^*|^{1-\gamma} \, dx - \lambda \|u^*\|_r^r \\
& + \varepsilon \int_{\Omega} \left( |\nabla u^*|^{p-2} \nabla u^* + \mu(x) |\nabla u^*|^{q-2} \nabla u^* \right) \cdot \nabla v \, dx \\
& - \varepsilon \int_{\Omega} \left( a(x) (u^*)^{-\gamma} + \lambda (u^*)^{r-1} \right) v \, dx \\
& - \int_{\{u^* + \varepsilon v < 0\}} \left( |\nabla u^*|^{p-2} \nabla u^* + \mu(x) |\nabla u^*|^{q-2} \nabla u^* \right) \cdot \nabla (u^* + \varepsilon v) \, dx \\
& + \int_{\{u^* + \varepsilon v < 0\}} \left( a(x) (u^*)^{-\gamma} + \lambda (u^*)^{r-1} \right) (u^* + \varepsilon v) \, dx \\
\leq & \varepsilon \int_{\Omega} \left( |\nabla u^*|^{p-2} \nabla u^* + \mu(x) |\nabla u^*|^{q-2} \nabla u^* \right) \cdot \nabla v \, dx \\
& - \varepsilon \int_{\Omega} \left( a(x) (u^*)^{-\gamma} + \lambda (u^*)^{r-1} \right) v \, dx \\
& - \varepsilon \int_{\{u^* + \varepsilon v < 0\}} \left( |\nabla u^*|^{p-2} \nabla u^* + \mu(x) |\nabla u^*|^{q-2} \nabla u^* \right) \cdot \nabla v \, dx.
\end{aligned}$$

Dividing the last inequality with  $\varepsilon > 0$  and letting  $\varepsilon \rightarrow 0$ , by taking

$$\int_{\{u^* + \varepsilon v < 0\}} \left( |\nabla u^*|^{p-2} \nabla u^* + \mu(x) |\nabla u^*|^{q-2} \nabla u^* \right) \cdot \nabla v \, dx \rightarrow 0 \quad \text{as } \varepsilon \rightarrow 0$$

into account, we obtain

$$\begin{aligned}
& \int_{\Omega} \left( |\nabla u^*|^{p-2} \nabla u^* + \mu(x) |\nabla u^*|^{q-2} \nabla u^* \right) \cdot \nabla v \, dx \\
& \geq \int_{\Omega} a(x) (u^*)^{-\gamma} v \, dx + \lambda \int_{\Omega} (u^*)^{r-1} v \, dx.
\end{aligned}$$

Since  $v \in W_0^{1,\mathcal{H}}(\Omega)$  is arbitrary chosen, equality must hold. It follows that  $u^*$  is a weak solution of problem (1.3) such that  $\varphi_{\lambda}(u^*) < 0$ , see Propositions 3.2 and 3.4.  $\square$

Now we start looking for a second weak solution when the parameter  $\lambda > 0$  is sufficiently small. To this end, we will use the manifold  $\mathcal{N}_{\lambda}^-$ .

**Proposition 3.8.** *Let hypotheses (H) be satisfied. Then there exists  $\hat{\lambda}_0^* \in (0, \hat{\lambda}^*]$  such that  $\varphi_{\lambda}|_{\mathcal{N}_{\lambda}^-} > 0$  for all  $\lambda \in (0, \hat{\lambda}_0^*]$ .*

*Proof.* From Proposition 3.4 we know that  $\mathcal{N}_{\lambda}^- \neq \emptyset$ . Let  $u \in \mathcal{N}_{\lambda}^-$ . By the definition of  $\mathcal{N}_{\lambda}^-$  and the embedding  $W_0^{1,p}(\Omega) \rightarrow L^r(\Omega)$  we have

$$\begin{aligned}
\lambda(r + \gamma - 1) \|u\|_r^r & > (p + \gamma - 1) \|\nabla u\|_p^p + (q + \gamma - 1) \|\nabla u\|_{q,\mu}^q \\
& \geq (p + \gamma - 1) \|\nabla u\|_p^p \\
& \geq (p + \gamma - 1) c_9^p \|u\|_r^p
\end{aligned}$$

for some  $c_9 > 0$ . Therefore

$$\|u\|_r \geq \left[ \frac{c_9^p (p + \gamma - 1)}{\lambda (r + \gamma - 1)} \right]^{\frac{1}{r-p}}. \quad (3.30)$$

Arguing by contradiction and suppose that the assertion of the proposition is not true. Then we can find  $u \in \mathcal{N}_\lambda^-$  such that  $\varphi_\lambda(u) \leq 0$ , that is,

$$\frac{1}{p} \|\nabla u\|_p^p + \frac{1}{q} \|\nabla u\|_{q,\mu}^q - \frac{1}{1-\gamma} \int_\Omega a(x)|u|^{1-\gamma} dx - \frac{\lambda}{r} \|u\|_r^r \leq 0. \quad (3.31)$$

Since  $\mathcal{N}_\lambda^- \subseteq \mathcal{N}_\lambda$  we know that

$$\frac{1}{q} \|\nabla u\|_{q,\mu}^q = \frac{1}{q} \int_\Omega a(x)|u|^{1-\gamma} dx + \frac{\lambda}{q} \|u\|_r^r - \frac{1}{q} \|\nabla u\|_p^p. \quad (3.32)$$

Using (3.32) in (3.31) gives

$$\left(\frac{1}{p} - \frac{1}{q}\right) \|\nabla u\|_p^p + \left(\frac{1}{q} - \frac{1}{1-\gamma}\right) \int_\Omega a(x)|u|^{1-\gamma} dx + \lambda \left(\frac{1}{q} - \frac{1}{r}\right) \|u\|_r^r \leq 0.$$

This yields

$$\lambda^{\frac{r-q}{qr}} \|u\|_r^r \leq \frac{q+\gamma-1}{q(1-\gamma)} \int_\Omega a(x)|u|^{1-\gamma} dx \leq \frac{q+\gamma-1}{q(1-\gamma)} c_{10} \|u\|_r^{1-\gamma}$$

for some  $c_{10} > 0$ . Therefore,

$$\|u\|_r \leq c_{11} \left(\frac{1}{\lambda}\right)^{\frac{1}{r+\gamma-1}} \quad (3.33)$$

for some  $c_{11} > 0$ . Now we use (3.33) in (3.30) in order to obtain

$$c_{12} \left(\frac{1}{\lambda}\right)^{\frac{1}{r-p}} \leq c_{11} \left(\frac{1}{\lambda}\right)^{\frac{1}{r+\gamma-1}} \quad \text{with} \quad c_{12} = \left[\frac{c_9^p(p+\gamma-1)}{r+\gamma-1}\right]^{\frac{1}{r-p}} > 0.$$

Consequently

$$0 < \frac{c_{12}}{c_{11}} \leq \lambda^{\frac{1}{r-p} - \frac{1}{r+\gamma-1}} = \lambda^{\frac{p+\gamma-1}{(r-p)(r+\gamma-1)}} \rightarrow 0 \quad \text{as} \quad \lambda \rightarrow 0^+,$$

since  $1 < p < r$  and  $\gamma \in (0, 1)$ , a contradiction. Thus, we can find  $\hat{\lambda}_0^* \in (0, \hat{\lambda}^*]$  such that  $\varphi_\lambda|_{\mathcal{N}_\lambda^-} > 0$  for all  $\lambda \in (0, \hat{\lambda}_0^*]$ .  $\square$

Now we minimize  $\varphi_\lambda$  on the manifold  $\mathcal{N}_\lambda^-$ .

**Proposition 3.9.** *Let hypotheses (H) be satisfied and let  $\lambda \in (0, \hat{\lambda}_0^*]$ . Then there exists  $v^* \in \mathcal{N}_\lambda^-$  with  $v^* \geq 0$  such that*

$$m_\lambda^- = \inf_{\mathcal{N}_\lambda^-} \varphi_\lambda = \varphi_\lambda(v^*) > 0.$$

*Proof.* Let  $\{v_n\}_{n \in \mathbb{N}} \subset \mathcal{N}_\lambda^- \subset \mathcal{N}_\lambda$  be a minimizing sequence. Since  $\mathcal{N}_\lambda^- \subset \mathcal{N}_\lambda$ , we know that  $\{v_n\}_{n \in \mathbb{N}} \subset W_0^{1,\mathcal{H}}(\Omega)$  is bounded, see Proposition 3.1. We may assume that

$$v_n \rightharpoonup v^* \quad \text{in} \quad W_0^{1,\mathcal{H}}(\Omega) \quad \text{and} \quad v_n \rightarrow v^* \quad \text{in} \quad L^r(\Omega).$$

Note that  $v^* \neq 0$  by (3.30). Now we will use the point  $t_2 > 0$  (see (3.12)) for which we have

$$\psi_{v^*}(t_2) = \lambda \|v^*\|_r^r \quad \text{and} \quad \psi'_{v^*}(t_2) < 0.$$

In the proof of Proposition 3.4 we showed that  $t_2 v^* \in \mathcal{N}_\lambda^-$ .

Next we want to show that  $\rho_{\mathcal{H}}(v_n) \rightarrow \rho_{\mathcal{H}}(v^*)$  as  $n \rightarrow \infty$  for a subsequence (still denoted by  $v_n$ ). Let us suppose this is not the case, then we have as in the proof of Proposition 3.4 that

$$\varphi_{\lambda}(t_2 v^*) < \lim_{n \rightarrow \infty} \varphi_{\lambda}(t_2 v_n).$$

We know that  $\varphi_{\lambda}(t_2 v_n) \leq \varphi_{\lambda}(v_n)$  since it is the global maximum because of  $\omega''_{v_n}(1) < 0$ . Using this and  $t_2 v^* \in \mathcal{N}_{\lambda}^-$ , we get

$$m_{\lambda}^- \leq \varphi_{\lambda}(t_2 v^*) < m_{\lambda}^-,$$

which is a contradiction. Hence we have  $\lim_{n \rightarrow +\infty} \rho_{\mathcal{H}}(v_n) = \rho_{\mathcal{H}}(v^*)$  for a subsequence and so Proposition 2.2(v) implies  $v_n \rightarrow v^*$  in  $W_0^{1,\mathcal{H}}(\Omega)$ . The continuity of  $\varphi_{\lambda}$  then gives  $\varphi_{\lambda}(v_n) \rightarrow \varphi_{\lambda}(v^*)$ , thus,  $\varphi_{\lambda}(v^*) = m_{\lambda}^-$ .

Since  $v_n \in \mathcal{N}_{\lambda}^-$  for all  $n \in \mathbb{N}$ , we have

$$(p + \gamma - 1) \|\nabla v_n\|_p^p + (q + \gamma - 1) \|\nabla v_n\|_{q,\mu}^q - \lambda(r + \gamma - 1) \|v_n\|_r^r < 0.$$

If we pass to the limit as  $n \rightarrow +\infty$  we obtain

$$(p + \gamma - 1) \|\nabla v^*\|_p^p + (q + \gamma - 1) \|\nabla v^*\|_{q,\mu}^q - \lambda(r + \gamma - 1) \|v^*\|_r^r \leq 0. \quad (3.34)$$

Recall that  $\lambda \in (0, \hat{\lambda}^*)$  and  $\hat{\lambda}^* \leq \lambda^*$ . Applying Proposition 3.3 we see that equality in (3.34) cannot happen. Hence,  $v^* \in \mathcal{N}_{\lambda}^-$ . Since the treatment also works for  $|v^*|$  instead of  $v^*$ , we may assume that  $v^*(x) \geq 0$  for a. a.  $x \in \Omega$  such that  $v^* \neq 0$ . Proposition 3.8 finally shows that  $m_{\lambda}^- > 0$ .  $\square$

Now we have a second weak solution of problem (1.3).

**Proposition 3.10.** *Let hypotheses (H) be satisfied and let  $\lambda \in (0, \hat{\lambda}_0^*]$ . Then  $v^*$  is a weak solution of problem (1.3) such that  $\varphi_{\lambda}(v^*) > 0$ .*

*Proof.* Following the proof of Proposition 3.6 replacing  $u^*$  by  $v^*$  in the definition of  $\eta_h$  by using Lemma 3.5 we are able to show that for every  $t \in [0, b_0]$  there exists  $\vartheta(t) > 0$  such that

$$\vartheta(t)(v^* + th) \in \mathcal{N}_{\lambda}^- \quad \text{and} \quad \vartheta(t) \rightarrow 1 \quad \text{as } t \rightarrow 0^+,$$

see also (3.25). Taking Proposition 3.9 into account we have that

$$m_{\lambda}^- = \varphi_{\lambda}(v^*) \leq \varphi_{\lambda}(\vartheta(t)(v^* + th)) \quad \text{for all } t \in [0, b_0]. \quad (3.35)$$

Let us now show that  $v^*(x) > 0$  for a. a.  $x \in \Omega$ . As for  $u^*$ , let us suppose there exists a set  $E$  with positive measure such that  $v^* = 0$  in  $E$ . Taking  $h \in W_0^{1,\mathcal{H}}(\Omega)$  with  $h > 0$  and  $t \in (0, b_0)$ , we know that  $(\vartheta(t)(v^* + th))^{1-\gamma} > (\vartheta(t)v^*)^{1-\gamma}$  a. e. in  $\Omega \setminus E$ . Note that  $\omega_{v^*}(1)$  is the global maximum which implies  $\omega_{v^*}(1) \geq \omega_{v^*}(\vartheta(t))$ . Using this and (3.35) it follows that

$$\begin{aligned} 0 &\leq \frac{\varphi_{\lambda}(\vartheta(t)(v^* + th)) - \varphi_{\lambda}(v^*)}{t} \\ &\leq \frac{\varphi_{\lambda}(\vartheta(t)(v^* + th)) - \varphi_{\lambda}(\vartheta(t)v^*)}{t} \\ &= \frac{1}{p} \frac{\|\nabla(\vartheta(t)(v^* + th))\|_p^p - \|\nabla(\vartheta(t)v^*)\|_p^p}{t} \\ &\quad + \frac{1}{q} \frac{\|\nabla(\vartheta(t)(v^* + th))\|_{q,\mu}^q - \|\nabla(\vartheta(t)v^*)\|_{q,\mu}^q}{t} - \frac{\vartheta(t)^{1-\gamma}}{(1-\gamma)t^{\gamma}} \int_E a(x)h^{1-\gamma} dx \end{aligned}$$

$$\begin{aligned}
& - \frac{1}{1-\gamma} \int_{\Omega \setminus E} a(x) \frac{(\vartheta(t)(v^* + th))^{1-\gamma} - (\vartheta(t)v^*)^{1-\gamma}}{t} dx \\
& - \frac{\lambda \|\vartheta(t)(v^* + th)\|_r^r - \|\vartheta(t)v^*\|_r^r}{r} \\
& < \frac{1}{p} \frac{\|\nabla(\vartheta(t)(v^* + th))\|_p^p - \|\nabla(\vartheta(t)v^*)\|_p^p}{t} \\
& + \frac{1}{q} \frac{\|\nabla(\vartheta(t)(v^* + th))\|_{q,\mu}^q - \|\nabla(\vartheta(t)v^*)\|_{q,\mu}^q}{t} - \frac{\vartheta(t)^{1-\gamma}}{(1-\gamma)t^\gamma} \int_E a(x) h^{1-\gamma} dx \\
& - \frac{\lambda \|\vartheta(t)(v^* + th)\|_r^r - \|\vartheta(t)v^*\|_r^r}{r}.
\end{aligned}$$

Therefore, similar to the proof of Proposition 3.7, we see from the inequality above that

$$0 \leq \frac{\varphi_\lambda(\vartheta(t)(v^* + th)) - \varphi_\lambda(\vartheta(t)v^*)}{t} \rightarrow -\infty \quad \text{as } t \rightarrow 0^+,$$

which is again a contradiction. We conclude that  $v^*(x) > 0$  for a. a.  $x \in \Omega$ .

The rest of the proof can be done similarly as the proof of Proposition 3.7. Precisely, (3.27) and (3.28) can be proven in the same way by applying again (3.35) and the inequality  $\omega_{v^*}(1) \geq \omega_{v^*}(\vartheta(t))$  together with  $v^* > 0$ . Finally, the last part of Proposition 3.7 is the same replacing  $u^*$  by  $v^*$ . From Proposition 3.9 we know that  $\varphi_\lambda(v^*) > 0$ . This finishes the proof.  $\square$

The proof of Theorem 1.1 follows now from Propositions 3.7 and 3.10.

#### ACKNOWLEDGMENT

The authors wish to thank Professor R. L. Alves for pointing out mistakes in the proofs of Propositions 3.6 and 3.7 in the first version of the manuscript.

W. Liu was supported by the NNSF of China (Grant No. 11961030). G. Dai is supported by the NNSF of China (Grant No. 11871129), the Fundamental Research Funds for the Central Universities (Grant No. DUT17LK05), and Xinghai Youqing funds from Dalian University of Technology.

#### REFERENCES

- [1] R.L. Alves, C.A. Santos, K. Silva, *Multiplicity of negative-energy solutions for singular-superlinear Schrödinger equations with indefinite-sign potential*, Commun. Contemp. Math., <https://doi.org/10.1142/S0219199721500425>.
- [2] A. Bahrouni, V.D. Rădulescu, P. Winkert, *Double phase problems with variable growth and convection for the Baouendi-Grushin operator*, Z. Angew. Math. Phys. **71** (2020), no. 6, 183, 14 pp.
- [3] G. Barletta, E. Tornatore, *Elliptic problems with convection terms in Orlicz spaces*, J. Math. Anal. Appl. **495** (2021), no. 2, 124779, 28 pp.
- [4] P. Baroni, M. Colombo, G. Mingione, *Harnack inequalities for double phase functionals*, Nonlinear Anal. **121** (2015), 206–222.
- [5] P. Baroni, M. Colombo, G. Mingione, *Non-autonomous functionals, borderline cases and related function classes*, St. Petersburg Math. J. **27** (2016), 347–379.
- [6] P. Baroni, M. Colombo, G. Mingione, *Regularity for general functionals with double phase*, Calc. Var. Partial Differential Equations **57** (2018), no. 2, Art. 62, 48 pp.
- [7] P. Baroni, T. Kuusi, G. Mingione, *Borderline gradient continuity of minima*, J. Fixed Point Theory Appl. **15** (2014), no. 2, 537–575.
- [8] S. Biagi, F. Esposito, E. Vecchi, *Symmetry and monotonicity of singular solutions of double phase problems*, J. Differential Equations **280** (2021), 435–463.

- [9] S.-S. Byun, J. Oh, *Global gradient estimates for non-uniformly elliptic equations*, Calc. Var. Partial Differential Equations **56** (2017), no. 2, Paper No. 46, 36 pp.
- [10] S.-S. Byun, J. Oh, *Regularity results for generalized double phase functionals*, Anal. PDE **13** (2020), no. 5, 1269–1300.
- [11] S.-S. Byun, Y. Youn, *Riesz potential estimates for a class of double phase problems*, J. Differential Equations **264** (2018), no. 2, 1263–1316.
- [12] Z.-Y. Chen, B. Ge, W.-S. Yuan, X.-F. Cao, *Existence of solution for double-phase problem with singular weights*, Adv. Math. Phys. **2020** (2020), Art. ID 5376013, 7 pp.
- [13] F. Colasuonno, M. Squassina, *Eigenvalues for double phase variational integrals*, Ann. Mat. Pura Appl. (4) **195** (2016), no. 6, 1917–1959.
- [14] M. Colombo, G. Mingione, *Bounded minimisers of double phase variational integrals*, Arch. Ration. Mech. Anal. **218** (2015), no. 1, 219–273.
- [15] M. Colombo, G. Mingione, *Calderón-Zygmund estimates and non-uniformly elliptic operators*, J. Funct. Anal. **270** (2016), no. 4, 1416–1478.
- [16] M. Colombo, G. Mingione, *Regularity for double phase variational problems*, Arch. Ration. Mech. Anal. **215** (2015), no. 2, 443–496.
- [17] Á. Crespo-Blanco, L. Gasiński, P. Harjulehto, P. Winkert, *A new class of double phase variable exponent problems: Existence and uniqueness*, J. Differential Equations, accepted 2022, <https://doi.org/10.1016/j.jde.2022.03.029>.
- [18] C. De Filippis, G. Mingione, *A borderline case of Calderón-Zygmund estimates for nonuniformly elliptic problems*, St. Petersburg Math. J. **31** (2020), no. 3, 455–477.
- [19] C. De Filippis, G. Mingione, *Lipschitz bounds and nonautonomous integrals*, Arch. Ration. Mech. Anal. **242** (2021), no. 2, 973–1057.
- [20] C. De Filippis, G. Mingione, *Manifold constrained non-uniformly elliptic problems*, J. Geom. Anal. **30** (2020), no. 2, 1661–1723.
- [21] C. De Filippis, G. Palatucci, *Hölder regularity for nonlocal double phase equations*, J. Differential Equations **267** (2019), no. 1, 547–586.
- [22] S. El Manouni, G. Marino, P. Winkert, *Existence results for double phase problems depending on Robin and Steklov eigenvalues for the  $p$ -Laplacian*, Adv. Nonlinear Anal. **11** (2022), no. 1, 304–320.
- [23] A. Esposito, F. Leonetti, G. Mingione, *Sharp regularity for functionals with  $(p, q)$  growth*, J. Differential Equations **204** (2004), no. 1, 5–55.
- [24] A. Esposito, F. Leonetti, P.V. Petricca *Absence of Lavrentiev gap for non-autonomous functionals with  $(p, q)$ -growth*, Adv. Nonlinear Anal. **8** (2019), no. 1, 73–78.
- [25] C. Farkas, P. Winkert, *An existence result for singular Finsler double phase problems*, J. Differential Equations **286** (2021), 455–473.
- [26] L. Gasiński, N.S. Papageorgiou, *Constant sign and nodal solutions for superlinear double phase problems*, Adv. Calc. Var. **14** (2021), no. 4, 613–626.
- [27] L. Gasiński, N.S. Papageorgiou, “Nonlinear Analysis”, Chapman & Hall/CRC, Boca Raton, FL, 2006.
- [28] L. Gasiński, P. Winkert, *Constant sign solutions for double phase problems with superlinear nonlinearity*, Nonlinear Anal. **195** (2020), 111739, 9 pp.
- [29] L. Gasiński, P. Winkert, *Existence and uniqueness results for double phase problems with convection term*, J. Differential Equations **268** (2020), no. 8, 4183–4193.
- [30] L. Gasiński, P. Winkert, *Sign changing solution for a double phase problem with nonlinear boundary condition via the Nehari manifold*, J. Differential Equations **274** (2021), 1037–1066.
- [31] P. Harjulehto, P. Hästö, “Orlicz Spaces and Generalized Orlicz Spaces”, Springer, Cham, 2019.
- [32] E. Hewitt, K. Stromberg, “Real and Abstract Analysis”, Springer-Verlag, New York, 1965.
- [33] C.-Y. Lei, *Existence and multiplicity of positive solutions for Neumann problems involving singularity and critical growth*, J. Math. Anal. Appl. **459** (2018), no. 2, 959–979.
- [34] W. Liu, G. Dai, *Existence and multiplicity results for double phase problem*, J. Differential Equations **265** (2018), no. 9, 4311–4334.
- [35] W. Liu, G. Dai, *Multiplicity results for double phase problems in  $\mathbb{R}^N$* , J. Math. Phys. **61** (2020), no. 9, 091508, 20 pp.
- [36] W. Liu, G. Dai, *Three ground state solutions for double phase problem*, J. Math. Phys. **59** (2018), no. 12, 121503, 7 pp.

- [37] P. Marcellini, *Regularity and existence of solutions of elliptic equations with  $p, q$ -growth conditions*, J. Differential Equations **90** (1991), no. 1, 1–30.
- [38] P. Marcellini, *Regularity of minimizers of integrals of the calculus of variations with non-standard growth conditions*, Arch. Rational Mech. Anal. **105** (1989), no. 3, 267–284.
- [39] P. Marcellini, *The stored-energy for some discontinuous deformations in nonlinear elasticity*, in “Partial differential equations and the calculus of variations, Vol. II”, vol. 2, 767–786, Birkhäuser Boston, Boston, 1989.
- [40] G. Marino, P. Winkert, *Existence and uniqueness of elliptic systems with double phase operators and convection terms*, J. Math. Anal. Appl. **492** (2020), 124423, 13 pp.
- [41] J. Musielak, “Orlicz Spaces and Modular Spaces”, Springer-Verlag, Berlin, 1983.
- [42] Z. Nehari, *Characteristic values associated with a class of non-linear second-order differential equations*, Acta Math. **105** (1961), 141–175.
- [43] Z. Nehari, *On a class of nonlinear second-order differential equations*, Trans. Amer. Math. Soc. **95** (1960), 101–123.
- [44] J. Ok, *Partial regularity for general systems of double phase type with continuous coefficients*, Nonlinear Anal. **177** (2018), 673–698.
- [45] J. Ok, *Regularity for double phase problems under additional integrability assumptions*, Nonlinear Anal. **194** (2020), 111408.
- [46] N.S. Papageorgiou, V.D. Rădulescu, D.D. Repovš, *Double-phase problems and a discontinuity property of the spectrum*, Proc. Amer. Math. Soc. **147** (2019), no. 7, 2899–2910.
- [47] N.S. Papageorgiou, V.D. Rădulescu, D.D. Repovš, *Existence and multiplicity of solutions for double-phase Robin problems*, Bull. Lond. Math. Soc. **52** (2020), no. 3, 546–560.
- [48] N.S. Papageorgiou, V.D. Rădulescu, D.D. Repovš, “Nonlinear Analysis – Theory and Methods”, Springer, Cham, 2019.
- [49] N.S. Papageorgiou, D.D. Repovš, C. Vetro, *Positive solutions for singular double phase problems*, J. Math. Anal. Appl. **501** (2021), no. 1, 123896, 13 pp.
- [50] N.S. Papageorgiou, C. Vetro, F. Vetro, *Solutions for parametric double phase Robin problems*, Asymptot. Anal. **121** (2021), no. 2, 159–170.
- [51] N.S. Papageorgiou, P. Winkert, *Positive solutions for weighted singular  $p$ -Laplace equations via Nehari manifolds*, Appl. Anal. **100** (2021), no. 11, 2436–2448.
- [52] K. Perera, M. Squassina, *Existence results for double-phase problems via Morse theory*, Commun. Contemp. Math. **20** (2018), no. 2, 1750023, 14 pp.
- [53] V.D. Rădulescu, *Isotropic and anisotropic double-phase problems: old and new*, Opuscula Math. **39** (2019), no. 2, 259–279.
- [54] M.A. Ragusa, A. Tachikawa, *Regularity for minimizers for functionals of double phase with variable exponents*, Adv. Nonlinear Anal. **9** (2020), no. 1, 710–728.
- [55] G. Riey, *Regularity and weak comparison principles for double phase quasilinear elliptic equations*, Discrete Contin. Dyn. Syst. **39** (2019), no. 8, 4863–4873.
- [56] Y. Sun, S. Wu, Y. Long, *Combined effects of singular and superlinear nonlinearities in some singular boundary value problems*, J. Differential Equations **176** (2001), no. 2, 511–531.
- [57] A. Szulkin, T. Weth, “The Method of Nehari Manifold”, Handbook of nonconvex analysis and applications, 597–632, Int. Press, Somerville, MA, 2010.
- [58] X. Wang, L. Zhao, P. Zhao, *Combined effects of singular and critical nonlinearities in elliptic problems*, Nonlinear Anal. **87** (2013), 1–10.
- [59] S. Zeng, Y. Bai, L. Gasiński, P. Winkert, *Convergence analysis for double phase obstacle problems with multivalued convection term*, Adv. Nonlinear Anal. **10** (2021), no. 1, 659–672.
- [60] S. Zeng, Y. Bai, L. Gasiński, P. Winkert, *Existence results for double phase implicit obstacle problems involving multivalued operators*, Calc. Var. Partial Differential Equations **59** (2020), no. 5, 176.
- [61] S. Zeng, L. Gasiński, P. Winkert, Y. Bai, *Existence of solutions for double phase obstacle problems with multivalued convection term*, J. Math. Anal. Appl. **501** (2021), no. 1, 123997, 12 pp.
- [62] V.V. Zhikov, *Averaging of functionals of the calculus of variations and elasticity theory*, Izv. Akad. Nauk SSSR Ser. Mat. **50** (1986), no. 4, 675–710.
- [63] V.V. Zhikov, S.M. Kozlov, O.A. Oleĭnik, “Homogenization of Differential Operators and Integral Functionals”, Springer-Verlag, Berlin, 1994.

(W. Liu) SCHOOL OF SCIENCE, JIANGXI UNIVERSITY OF SCIENCE AND TECHNOLOGY, GANZHOU,  
JIANGXI 341000, PR CHINA

*Email address:* liuwul000@gmail.com

(G. Dai) SCHOOL OF MATHEMATICAL SCIENCES, DALIAN UNIVERSITY OF TECHNOLOGY, DALIAN,  
116024, PR CHINA

*Email address:* daiguowei@dlut.edu.cn

(N. S. Papageorgiou) NATIONAL TECHNICAL UNIVERSITY, DEPARTMENT OF MATHEMATICS, ZO-  
GRAFOU CAMPUS, ATHENS 15780, GREECE

*Email address:* npapg@math.ntua.gr

(P. Winkert) TECHNISCHE UNIVERSITÄT BERLIN, INSTITUT FÜR MATHEMATIK, STRASSE DES  
17. JUNI 136, 10623 BERLIN, GERMANY

*Email address:* winkert@math.tu-berlin.de